



كلية الطب  
والصيدلة - مراكش  
FACULTÉ DE MÉDECINE  
ET DE PHARMACIE - MARRAKECH

Year : 2019Thesis N°234

## Adhesive small bowel obstruction in children

### THESIS

PRESENTED AND DEFENDED PUBLICLY THE 01/10/2019

BY

**Mr. YOUNESS EL-KHADIR**

Born on the 27 September 1993 in Kasba Tadla

TO OBTAIN THE DEGREE OF DOCTOR OF MEDECINE

### KEYWORDS :

Adhesions – Adhesive small bowel obstruction – ASBO – Children – Pediatric – conservative management – Water-soluble contrast – Gastrographin

### JURY

Mr. **S. YOUNOUS**

Professor of Anesthesiology and Intensive Care Medicine

PRESIDENT

Mr. **M. OULAD SAIAD**

Professor of General Pediatric Surgery

SUPERVISOR

Mr. **E.E. KAMILI**

Professor of General Pediatric Surgery

Mr. **H. JALAL**

Professor of Radiology

Mr. **M. BOURROUS**

Professor of Pediatrics

JUDGES

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَقَدْ عَلِمْتُمُ

# Hippocratic Oath

*I swear to fulfill, to the best of my ability and judgment, this covenant:*

*I will respect the hard-won scientific gains of those physicians in whose steps I walk, and gladly share such knowledge as is mine with those who are to follow.*

*I will apply, for the benefit of the sick, all measures [that] are required, avoiding those twin traps of overtreatment and therapeutic nihilism.*

*I will remember that there is art to medicine as well as science, and that warmth, sympathy, and understanding may outweigh the surgeon's knife or the chemist's drug.*

*I will not be ashamed to say "I know not," nor will I fail to call in my colleagues when the skills of another are needed for a patient's recovery.*

*I will respect the privacy of my patients, for their problems are not disclosed to me that the world may know. Most especially must I tread with care in matters of life and death. If it is given me to save a life, all thanks. But it may also be within my power to take a life; this awesome responsibility must be faced with great humbleness and awareness of my own frailty. Above all, I must not play at God.*

*I will remember that I do not treat a fever chart, a cancerous growth, but a sick human being, whose illness may affect the person's family and economic stability. My responsibility includes these related problems, if I am to care adequately for the sick.*


*I will prevent disease whenever I can, for prevention is preferable to cure.*

*I will remember that I remain a member of society, with special obligations to all my fellow human beings, those sound of mind and body as well as the infirm.*

*If I do not violate this oath, may I enjoy life and art, respected while I live and remembered with affection thereafter. May I always act so as to preserve the finest traditions of my calling and may I long experience the joy of healing those who seek my help.*

**Declaration of Geneva, 1948**





---

*LIST OF PROFESSORS*

---



**UNIVERSITE CADI AYYAD**  
**FACULTE DE MEDECINE ET DE PHARMACIE**  
**MARRAKECH**

Doyens Honoraires

: Pr. Badie Azzaman MEHADJI  
: Pr. Abdelhaq ALAOUI YAZIDI

ADMINISTRATION

Doyen

: Pr. Mohammed BOUSKRAOUI

Vice doyen à la Recherche et la Coopération

: Pr. Mohamed AMINE

Vice doyen aux Affaires Pédagogiques

: Pr. Redouane EL FEZZAZI

Secrétaire Générale

: Mr. Azzeddine ELHOUDAIGUI

**Professeurs de  
l'enseignement supérieur**

Nom et Prénom	Spécialité	Nom et Prénom	Spécialité
ABKARI Imad	Traumato- orthopédie	FINECH Benasser	Chirurgie – générale
ABOU EL HASSAN Taoufik	Anesthésie- réanimation	FOURAIJI Karima	Chirurgie pédiatrique
ABOUCHADI Abdeljalil	Stomatologie et chir maxillo faciale	GHANNANE Houssine	Neurochirurgie
ABOULFALAH Abderrahim	Gynécologie- obstétrique	GHOUNDALE Omar	Urologie
ABOUSSAIR Nisrine	Génétique	HAJJI Ibtissam	Ophtalmologie
ADERDOUR Lahcen	Oto- rhino- laryngologie	HOCAR Ouafa	Dermatologie
ADMOU Brahim	Immunologie	JALAL Hicham	Radiologie
AGHOUTANE El Mouhtadi	Chirurgie pédiatrique	KAMILI El Ouafi El Aouni	Chirurgie pédiatrique
AIT AMEUR Mustapha	Hématologie Biologique	KHALLOUKI Mohammed	Anesthésie- réanimation
AIT BENALI Said	Neurochirurgie	KHATOURI Ali	Cardiologie

AIT BENKADDOURYassir	Gynécologie- obstétrique	KHOUCHANIMouna	Radiothérapie
AIT-SABImane	Pédiatrie	KISSANINajib	Neurologie
AKHDARINadia	Dermatologie	KOULALI IDRISIKhalid	Traumato-orthopédie
ALAOUIMustapha	Chirurgie- vasculaire périphérique	KRATIKhadija	Gastro-entérologie
AMALSaid	Dermatologie	KRIETMohamed	Ophtalmologie
AMINEMohamed	Epidémiologie- clinique	LAGHMARIMehdi	Neurochirurgie
AMMAR Haddou	Oto-rhino-laryngologie	LAKMICHIMohamed Amine	Urologie
AMROLamyae	Pneumo- phtisiologie	LAOUADInass	Néphrologie
ARSALANELamiaie	Microbiologie -Virologie	LOUZIAbdelouahed	Chirurgie – générale
ASMOUKIHamid	Gynécologie- obstétrique	MADHAR Si Mohamed	Traumato-orthopédie
ASRIFatima	Psychiatrie	MANOUDIFatiha	Psychiatrie
BEN DRISSLaila	Cardiologie	MANSOURINadia	Stomatologie et chiru maxillo faciale
BENCHAMKHAYassine	Chirurgie réparatrice et plastique	MOUDOUNISaid Mohammed	Urologie
BENELKHAIAT BENOMARRidouan	Chirurgie - générale	MOUFIDKamal	Urologie
BENJILALILaila	Médecineinterne	MOUTAJ Redouane	Parasitologie
BOUAITY Brahim	Oto-rhino- laryngologie	MOUTAOUAKILAbdeljalil	Ophtalmologie
BOUCHENTOUFRachid	Pneumo- phtisiologie	NAJEBYoussef	Traumato-orthopédie
BOUGHALEM Mohamed	Anesthésie - réanimation	NARJISSYoussef	Chirurgiegénérale
BOUKHIRAAbderrahman	Biochimie - chimie	NEJMI Hicham	Anesthésie-réanimation
BOUMZEBRADrissi	Chirurgie Cardio-Vasculaire	NIAMANE Radouane	Rhumatologie
BOURROUSMonir	Pédiatrie	NOURIHassan	Oto rhino laryngologie
BOUSKRAOUMohammed	Pédiatrie	OUALI IDRISIMariem	Radiologie
CHAFIKRachid	Traumato- orthopédie	OULAD SAIADMohamed	Chirurgie pédiatrique
CHAKOUR Mohamed	Hématologie Biologique	QACIF Hassan	Médecineinterne
CHELLAKSaliha	Biochimie- chimie	QAMOUSSYoussef	Anesthésie-réanimation

CHERIF IDRISSE EL GANOUNI Najat	Radiologie	RABBANIKhalid	Chirurgie générale
CHOULLI MohamedKhaled	Neuro pharmacologie	RAFIK Redda	Neurologie
DAHAMI Zakaria	Urologie	RAJIAbdelaziz	Oto-rhino-laryngologie
EL ADIB Ahmed Rhassane	Anesthésie- réanimation	SAIDI Halim	Traumatologie- orthopédie
EL ANSARINawal	Endocrinologie et maladies métaboliques	SAMKAOUI Mohamed Abdenasser	Anesthésie- réanimation
EL BARNIRachid	Chirurgie- générale	SAMLANI Zouhour	Gastro- entérologie
EL BOUCHTIImane	Rhumatologie	SARFIsmail	Urologie
EL BOUIHIMohamed	Stomatologie et chir maxillo faciale	SORANabila	Microbiologie - Virologie
ELFEZZAZI Redouane	Chirurgie pédiatrique	SOUMMANIAbderraouf	Gynécologie- obstétrique
EL HAOURYHanane	Traumatologie- orthopédie	TASSINoura	Maladies infectieuses
EL HATTAOUMustapha	Cardiologie	YOUNOUSSaid	Anesthésie- réanimation
EL HOUDZIJamila	Pédiatrie	ZAHLANEMouna	Médecine interne
EL KARIMISaloua	Cardiologie	ZOUHAIR Said	Microbiologie
ELFIKRIAbdelghani	Radiologie	ZYANI Mohammed	Médecine interne
ESSAADOUNILamiaa	Médecine interne		

### Professeurs Agrégés

Nom et Prénom	Spécialité	Nom et Prénom	Spécialité
ABIR Badreddine	Stomatologie et Chirurgie maxillo faciale	GHAZI Miriema	Rhumatologie
ADALIImane	Psychiatrie	HACHIMIAbdelhamid	Réanimation médicale
ADARMOUCH Latifa	Médecine Communautaire (médecine préventive, santé publique et hygiène)	HAROUKaram	Gynécologie- obstétrique
AISSAOUIYounes	Anesthésie - réanimation	HAZMIRI Fatima Ezzahra	Histologie – Embryologie - Cytogénétique
AIT BATAHAR Salma	Pneumo- phtisiologie	IHBIBANE fatima	Maladies Infectieuses

ALJSoumaya	Radiologie	KADDOURI Said	Médecineinterne
ANIBAKhalid	Neurochirurgie	LAHKIM Mohammed	Chirurgiegénérale
ATMANE El Mehdi	Radiologie	LAKOUICHMIMohammed	Stomatologie et Chirurgie maxillo faciale
BAIZRIHicham	Endocrinologie et maladies métaboliques	LOUHABNisrine	Neurologie
BASRAOUIDounia	Radiologie	MAOULAININE Fadl mrabih rabou	Pédiatrie (Neonatalogie)
BASSIRAhlam	Gynécologie- obstétrique	MARGAD Omar	Traumatologie - orthopédie
BELBACHIR Anass	Anatomie- pathologique	MATRANEAboubakr	Médecinenucléair e
BELBARAKARhizlane	Oncologiemédicale	MEJDANEAbdelhadi	Chirurgie Générale
BELKHOUAhlam	Rhumatologie	MLIHA TOUATI Mohammed	Oto-Rhino - Laryngologie
BENHIMA Mohamed Amine	Traumatologie - orthopédie	MOUAFFAKYoussef	Anesthésie - réanimation
BENJELLOUN HARZIMI Amine	Pneumo- phtisiologie	MOUHSINE Abdelilah	Radiologie
BENLAIAbdeslam	Psychiatrie	MSOUGGARYassine	Chirurgiethoraciq ue
BENZAROUELDounia	Cardiologie	NADER Youssef	Traumatologie - orthopédie
BOUKHANNILahcen	Gynécologie- obstétrique	OUBAHA Sofia	Physiologie
BOURRAHOUATAicha	Pédiatrie	RADANoureddine	Pédiatrie
BSISS Mohamed Aziz	Biophysique	RAISHanane	Anatomiepatholo gique
CHRAA Mohamed	Physiologie	RBAIBI Aziz	Cardiologie
DAROUASSIYoussef	Oto-Rhino - Laryngologie	ROCHDIYoussef	Oto-rhino- laryngologie
DRAISSGhizlane	Pédiatrie	SAJIAIHafsa	Pneumo- phtisiologie
EL AMRANI MoulayDriss	Anatomie	SALAMATarik	Chirurgiepédiatriq ue
EL HAOUATIRachid	Chirurgie Cardio- vasculaire	SEDDIKI Rachid	Anesthésie - Réanimation
EL IDRISSE SLITINENadia	Pédiatrie	SERGHINI Issam	Anesthésie - Réanimation
EL KHADER Ahmed	Chirurgiegénérale	TAZI Mohamed Illias	Hématologie- clinique
EL KHAYARIMina	Réanimationmédicale	TOURABI Khalid	Chirurgie réparatrice et plastique


EL MEZOUARI El Moustafa	Parasitologie Mycologie	ZAHLANE Kawtar	Microbiologie - virologie
EL MGHARI TABIB Ghizlane	Endocrinologie et maladies métaboliques	ZAOUISanaa	Pharmacologie
EL OMRANI Abdelhamid	Radiothérapie	ZARROUKI Youssef	Anesthésie - Réanimation
FADILI Wafaa	Néphrologie	ZEMRAOUI Nadir	Néphrologie
FAKHIR Bouchra	Gynécologie- obstétrique	ZIADI Amra	Anesthésie - réanimation
FAKHRI Anass	Histologie- embryologie cytogénétique	ZIDANE Moulay Abdelfettah	Chirurgie Thoracique

### Professeurs Assistants

Nom et Prénom	Spécialité	Nom et Prénom	Spécialité
ABDEL FETTAH Youness	Rééducation et Réhabilitation Fonctionnelle	ELOUARDI Youssef	Anesthésie réanimation
ABDOU Abdessamad	Chiru Cardio vasculaire	ELQATNI Mohamed	Médecine interne
AIT ERRAMI Adil	Gastro-entérologie	ESSADI Ismail	Oncologie Médicale
AKKA Rachid	Gastro - entérologie	FDIL Naima	Chimie de Coordination Bio-organique
ALAOUI Hassan	Anesthésie - Réanimation	FENNANE Hicham	Chirurgie Thoracique
AMINE Abdellah	Cardiologie	GHOZLANI Imad	Rhumatologie
ARABI Hafid	Médecine physique et réadaptation fonctionnelle	HAJJIFouad	Urologie
ARSALANE Adil	Chirurgie Thoracique	HAMMI Salah Eddine	Médecine interne
ASSERRAJI Mohammed	Néphrologie	Hammoune Nabil	Radiologie
AZIZ Zakaria	Stomatologie et chirurgie maxillo faciale	JALLAL Hamid	Cardiologie
BAALLAL Hassan	Neurochirurgie	JANAH Hicham	Pneumo-phtisiologie
BABA Hicham	Chirurgie générale	LAFFINTI Mahmoud Amine	Psychiatrie
BELARBI Marouane	Néphrologie	LAHLIMI Fatima Ezzahra	Hématologie clinique
BELFQUIH Hatim	Neurochirurgie	LALYA Issam	Radiothérapie
BELGHMAIDI Sarah	OPhtalmologie	LOQMAN Souad	Microbiologie et toxicologie environnementale
BELHADJ Ayoub	Anesthésie - Réanimation	MAHFOUD Tarik	Oncologie médicale

BELLASRI Salah	Radiologie	MILOUDI Mohcine	Microbiologie - Virologie
BENANTAR Lamia	Neurochirurgie	MOUNACH Aziza	Rhumatologie
BENNAOUI Fatiha	Pédiatrie	NAOUI Hafida	Parasitologie Mycologie
BOUCHENTOUF Sidi Mohammed	Chirurgie générale	NASSIH Houda	Pédiatrie
BOUKHRIS Jalal	Traumatologie - orthopédie	NASSIM SABAH Taoufik	Chirurgie Réparatrice et Plastique
BOUTAKIOUTE Badr	Radiologie	NYA Fouad	Chirurgie Cardio - Vasculaire
BOUZERDA Abdelmajid	Cardiologie	OUEIAGLI NABIH Fadoua	Psychiatrie
CHETOUI Abdelkhalek	Cardiologie	OUMERZOUK Jawad	Neurologie
CHETTATI Mariam	Néphrologie	RAISSI Abderrahim	Hématologie clinique
DAMI Abdallah	Médecine Légale	REBAHI Houssam	Anesthésie - Réanimation
DOUIREK Fouzia	Anesthésie- réanimation	RHARRASSI Isam	Anatomie- patologique
EL- AKHIRI Mohammed	Oto- rhino- laryngologie	SAOUAB Rachida	Radiologie
EL AMIRI My Ahmed	Chimie de Coordination bio-organique	SAYAGH Sanae	Hématologie
EL FAKIRI Karima	Pédiatrie	SEBBANI Majda	Médecine Communautaire (médecine préventive, santé publique et hygiène)
EL HAKKOUNI Awatif	Parasitologie mycologie	TAMZAOURTE Mouna	Gastro - entérologie
EL HAMZAOUI Hamza	Anesthésie réanimation	WARDA Karima	Microbiologie
EL KAMOUNI Youssef	Microbiologie Virologie	ZBITOU Mohamed Anas	Cardiologie
ELBAZ Meriem	Pédiatrie	ELOUARDI Youssef	Anesthésie réanimation

**LISTE ARRÊTÉE LE**  
**22/04/2019**



---

# *DEDECATIONS*

---



# بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

"رَبِّ أَوْزِعْنِي أَنْ أَشْكُرَ نِعْمَتَكَ الَّتِي

أَنْعَمْتَ عَلَيَّ وَعَلَىٰ وَالِدَيَّ وَأَنْ أَعْمَلَ

صَالِحًا تَرْضَاهُ وَأَدْخِلْنِي بِرَحْمَتِكَ فِي

عِبَادِكَ الصَّالِحِينَ."

صدق الله العظيم

سورة النمل الآية 19

*First and foremost praise is to ALLAH, the Almighty, the Greatest of all, on whom ultimately we depend for sustenance and guidance.*

*I would like to thank Almighty ALLAH for giving me the opportunity, determination and strength to complete my thesis. His continuous grace and mercy were with me throughout my life.*

*To the beloved memory of my Father Mohamed EL-KHADIR  
(1958-2019), may ALLAH bless his soul and grant him the  
highest levels of Jannah.*

*I could write a million words, but still be unable to say just how  
much I love and miss him every single day.*

*I will always remember him as a Superhero, a kind and loving father  
who had a heart of gold .and I will never forget all he taught me.  
His absence hurt me, but I won't be sad, because I know for sure  
that he is in better place, watching me and proud of me. I will  
always hold his memory in my mind and my heart.*

*To my beloved mother, NAJIA ASSAKAIE.*

*All the existing words of love in this universe can't express the endless gratitude and the love that I carry for you. If I could write a story, it would be the best story of a brave mother who protects and provides for her children, a perfect mother with the most beautiful soul and the purest heart on this earth .every step I took was made easier with your blessed prayers and your supports. You have always been, my source of tenderness, my inspiration, my guidance that whispers wisdom in my ears and enlightens my path.*

*My dear mom, I owe you everything in my life, and I hope that I have been up to your expectations and I have made you so proud as you have always been my source of pride and my example. May Almighty ALLAH protect you and grant you health, long life, and happiness.*

*I LOVE YOU SO MUCH MOM.*

*To my dears beautiful sisters LEILA, HIND, and  
MAROUA*

*It is not a lie when they said that there is nothing more beautiful in this life than having sisters. During my whole life you have been my angels that support me and give me strength. You are the most beautiful, brave, and kind sisters that everyone dreams to have. I feel very lucky and proud of you and I hope I have been a good brother for you and a source of pride. You are always in my heart my beloved angels. I wish you all the best and may ALLAH bless and protect you and guide your steps.*

*To my dearest aunts HALIMA and AICHA*

*Since my birth and for all those years You have been a second moms for me, you raised me and you looked after me , you showed me affection and tenderness. I think that all the words of love and gratitude wont be enough to express how grateful I am to you . I love you my Moms and I thank you for everything . I ask ALLAH to protect you ,and grant you health , long life and happiness .*

*To my beloved little birds, my niece Rayhane Sakhi and my  
nephew Mohamed Reda Sakhi*

*Of the many blessings in my life, I count you as a main one. If  
nephews and nieces were jewels, I would have the most beautiful  
gems ever. I am very lucky and proud of you. You have brought for  
me and for all the family joy and happiness. I ask ALLAH to  
protect you, May you have a wonderful year ahead of you and may  
all your dreams come true.*

*To the memory of my beloved Grandmothers Rabha  
Moustabchir and Fettouma El boukhari and my Grandfather  
Moha El-khadir*

*Your special love has always submerged my life. Thank you for your  
sincere prayers that helped me through life. I pray Allah to grant you  
the highest ranks of Jannah.*

*To all my adorable family members :*

*My Grandfather Ibrahim ASSAKAIE,*

*My Uncles and Aunts,*

*Mr Othmane Sakhi,*

*my cousins Hasna, Mehdi, Khaled, Badr, Imane, Aya,  
Hidaya, Manal, fakhreedine, Tawab and Sojoud*

*my childhood best friends and brothers: Abdelkadir Douali,  
Marouane Intissar, Hakim El Amrani, Ayoub Aguelmam and  
Hamza Recharg.*

*Families are the first in our lives, as we grow we always returning  
to those who love, encourage, and support us the most, our family.  
Throughout the years I had many challenges, but you were still there  
offering support and encouragement. Your kindness, your generosity,  
and your help have been an inspiration for me to overcome all those  
challenges, Please find in this modest work the expression of my deep  
affection and my sincere gratitude*

*To all my best friends and colleagues:*

*My brothers ;Younes, Youssef, Abdellah, Mustapha, Zakaria,  
Walid, Samir, Salah, Othmane, Mohamed, Youness,  
Abderrahim, Hamza, Adnane, Abdelkarim, Abdelhamid,  
Anouar, Abdelali ,Mouad, Tawfiq, Abdelwahed, redoine, Imad,  
Ayoub ...*

*My sisters; Fadoua, Fatimezzahra, Mounia, Nisserine,  
Soukaina, Rania, Laila, Rihane, Sara, Nassima, jihade,  
Naima, Hajar, Joumana, Asma, Yasmine, Rokaya, Ichrak, Hanae,  
Majda ....*

*Throughout those past years we shared memorable moments We  
worked together, we laughed and cried together, we made up and  
shared many more precious moments of life. Our friendship  
overpassed the university benches and hospital wards, and it  
continues to bring me joy and happiness in my life.*

*To all of you, my friends that I have mentioned and my friends that  
I forgot to mention unintentionally. I am honored and so grateful  
that I had the opportunity to meet you. Thank you for giving me  
memories that I am going to hold close to my heart and cherish  
forever . and May you shine in your career and your personal lives*


*To all medical and paramedical staff of the Mohamed VI teaching Hospital of Marrakech , Hassan II hospital of Khouribga , & Moulay Ismail hospital of Kasba Tadla.*

*To all the people who have supported me during these years of medical studies.*

*To Sci-hub community*

*To all medical students and doctors of the Stethoscope Revolution*

*Please find this work, the testimony of my sincere gratitude ,and my deep appreciation and respect.*



---

# *ACKNOWLEDGMENTS*

---



*To my Dear Master and thesis president, Professor Said YOUNOUS, Professor of Anesthesiology and Intensive care medicine*

*Thank you for granting me this great honor by accepting the presidency of this honorable jury. You are the example of the professor with great human and professional qualities .your seriousness, your competence, and your sense of duty have always been a source of inspiration for me and all your students. Please accept through this work the expression of my sincere gratitude and my deep consideration and respect.*

*To my Dear Master and Thesis supervisor, Professor Mohamed OULAD SAIAD Professor of General Pediatric Surgery*

*I would like to express my sincere gratitude and my deep respect for trusting me to Conduct this study. Your kindness, your human and professional qualities deserve all of admiration, and they make of you a role model for me and for all your students and trainees. Thank you for your patience, your pertinent advices, and for your guidance in every step during this work, I owe you this research experience, and I hope that I have been up to your expectations.*

*To my Dear Master and thesis judge, Professor El Ouafi El Aouni  
KAMILI Professor of General Pediatric Surgery*

*I would like to express my sincere gratitude and my deep consideration for your valuable participation and help in the development of this work and for your tireless encouragement. Your kindness, your generosity and your humility that you share your knowledge with, deserves all of respect and esteem. thank you for accepting to be part of this jury .and please find here, the testimony of my high consideration and deep appreciation*

*To my Dear Master and thesis judge, Professor Mounir  
BOURROUS , professor of Pediatrics*

*I sincerely thank you for granting me the honor by accepting to be part of this honorable committee. I always appreciated your kindness, your humility and your competence that make of you a great Master that every medical student and trainee doctor dream to have. Please find this work, the testimony of my sincere gratitude and my high esteem and respect*

*To my Dear Master and thesis judge, Professor Hicham JALAL ,  
Professor of Radiology*

*You have granted me a great honor by accepting being part of this respectful jury. The kindness you have shown while receiving this thesis was particularly touching. I thank you for your availability, your kindness and your professional dedication that make you a great Master to follow. please find here the expression of my sincere gratitude and my deep consideration*

## **List of abbreviations :**

ASBO : Adhesive small bowel obstruction

AXR : Abdominal X-ray

NOM: Non operative management

AFL : air-fluid levels

DLB : Dilated loops of bowel

CT scan : Computed tomography scan

WSCA : Water-soluble contrast agent

CRP : C-Reactive Protein

BUN : Blood urea nitrogen

LDH :Lactate dehydrogenase

CK : Creatine kinase

OIB : Occlusion intestinale par brides



---


# *TABLE OF CONTENTS*

---



<b>INTRODUCTION</b> .....	-1-
<b>PATIENTS &amp; METHODS</b> .....	-3-
<b>I. TYPE OF STUDY</b> .....	-4-
<b>II. PURPOSE OF THE STUDY</b> .....	-4-
<b>III. PATIENTS</b> .....	-4-
1. Inclusion criteria.....	-4-
2. Exclusion criteria .....	-4-
<b>IV. METHODS</b> .....	-5-
<b>RESULTS</b> .....	-6-
<b>I. EPIDEMIOLOGICAL DATA</b> .....	-7-
1. Incidence .....	-7-
2. Gender.....	-7-
3. Age .....	-8-
<b>II. PAST MEDICAL HISTORY</b> .....	-8-
1. Medical history .....	-8-
2. Surgical history .....	-8-
<b>III. HISTORY OF PRESENTING ILLNESS</b> .....	-10-
1. Symptoms .....	-10-
2. the onset of symptoms .....	-10-
3. Patient delay.....	-11-
4. the Time period between the previous surgery and development of ASBO symptoms .....	-11-
<b>IV. PHYSICAL EXAMINATION</b> .....	-12-
1. General examination.....	-12-
2. Abdominal examination .....	-12-
<b>V. PARACLINICAL INVESTIGATIONS</b> .....	-13-
1. RADIOLOGICAL INVESTIGATIONS.....	-13-
2. BIOLOGICAL TESTS .....	-15-
<b>VI. MANAGEMENT</b> .....	-16-
1. Conservative management .....	-16-
2. Operative management.....	-19-
<b>VII. EVOLUTION</b> .....	-20-
1. Length of hospital stay .....	-20-
2. Recurrence.....	-21-
3. Mortality .....	-21-

<b>DISCUSSION</b> .....	<b>-22-</b>
<b>I. EPIDEMIOLOGICAL DATA</b> .....	<b>-23-</b>
1. Gender.....	-23-
2. Age .....	-24-
3. PAST MEDICAL HISTORY .....	-25-
<b>II. PATHOPHYSIOLOGY</b> .....	<b>-29-</b>
1. Etiopathogenesis of adhesions .....	-29-
2. Mechanisms of obstruction.....	-33-
3. Consequences of intestinal obstruction.....	-33-
<b>III. CLINICAL FEATURES</b> .....	<b>-37-</b>
1. Symptoms.....	-37-
2. Physical signs .....	-38-
<b>IV. PARACLINICAL FEATURES</b> .....	<b>-39-</b>
1. Radiological investigations .....	-39-
2. Biological tests .....	-47-
<b>V. MANAGEMENT</b> .....	<b>-48-</b>
1. The Goals of treatment .....	-48-
2. Therapeutic approaches .....	-48-
2.1. Conservative management.....	-48-
2.2. Operative management.....	-55-
<b>VI. EVOLUTION</b> .....	<b>-64-</b>
1. Length of hospital stay .....	-64-
2. Recurrence.....	-65-
3. Mortality .....	-66-
<b>VIII. PREVENTION</b> .....	<b>-67-</b>
<b>CONCLUSION</b> .....	<b>-70-</b>
<b>APPENDIX</b> .....	<b>-73-</b>
<b>ABSTRACT</b> .....	<b>-85-</b>
<b>BIBLIOGRAPHY</b> .....	<b>-89-</b>



---

# *INTRODUCTION*

---



Adhesive small bowel obstruction (ASBO) is a mechanical disruption in the patency of the gastrointestinal tract, caused by peritoneal adhesions and resulting in a combination of emesis, obstipation, and abdominal pain.


Peritoneal adhesions can be defined as abnormal fibrous bands between organs or tissues or both in the abdominal cavity that are usually separated. They may be acquired or congenital. However, most are acquired as a result of abdominopelvic surgery, Less commonly, adhesions may form as the result of inflammatory conditions, intraperitoneal infection, radiation, or abdominal trauma [1]

Adhesive small bowel obstruction (ASBO) is very common in adults and considered the second cause of intestinal obstruction after obstructed abdominal wall hernia [2]. Moreover, it was reported that up to 25% of adult patients who have undergone abdominal surgery, subsequently developed ASBO [3]. Unfortunately, this condition does not spare children; the available data reported an incidence of ASBO in children that range between 1% and 9% [4–7].

ASBO should be suspected in any child with surgical history, who is presenting persistent vomiting, distension, and abdominal pain. Moreover, further investigations must be performed to confirm the diagnosis because delayed diagnosis and treatment can lead to devastating consequences. [8]

In adult patients with ASBO, conservative management found to be effective in a large proportion of cases [10–12], as well as operative management which can be effective and even essential in some cases, though it carries a risk of associated morbidity and mortality[9] . in children , there is a lack of consensus, or available guidelines, on the management of ASBO ; furthermore a small number of existing studies in the broader literature that have reviewed those therapeutic approaches in this group age .

The present study was designed to review the epidemiological, the clinical, and the paraclinical features, and mainly, to outline and compare the different aspects of ASBO management in children.



---

## *PATIENTS & METHODS*

---



## **I. TYPE OF STUDY :**

This is a retrospective study, that involved 61 cases of adhesive small bowel obstruction (ASBO) in 57 children admitted in the pediatric surgery department of the Mohamed VI Teaching Hospital of Marrakech, within a period of 7 years, between January 2012 and December 2018

## **II. PURPOSE OF THE STUDY**

This study aimed to review the epidemiological, the clinical, and the paraclinical features in children with ASBO .and also to outline and compare the existing therapeutic approaches for managing ASBO in pediatric patients .( conservative management Vs. operative management), and review their outcomes

## **III. PATIENTS**

### **1. INCLUSION CRITERIA**

Our study involved children under 15 years old, who were admitted in the pediatric surgery department of the Mohamed VI Teaching Hospital of Marrakech, and they were diagnosed with adhesive small bowel obstruction, basing on the past medical history, the clinical presentation and the abdominal X-ray

### **2. EXCLUSION CRITERIA**

Patients diagnosed with other causes of small bowel obstruction were excluded.

## IV. METHODS

to make this work, we followed those steps :

❖ Step1 :

Using the registry of our pediatric surgery department, we searched for medical records of children with ASBO that fit our criteria

❖ Step2 :

We made a medical summary sheet (see appendix I ) which contains different parts for our studied parameters including ; Identity, past medical history, history of the present complain, clinical and preclinical features, management, outcome, and evolution of the patient.

❖ Step 3 :

we collected the essential data from medical records then we gathered them in the summary medical sheets.

❖ step 4 :

the gathered data were entered into an Excel database then organized and analyzed



# *RESULTS*



## I. EPIDEMIOLOGICAL DATA

### 1. Incidence :


During a 7 years period, between January 2012 and December 2018. We have identified 61 episodes of ASBO in 57 patients admitted in our pediatric surgery department.

**Table I:** Distribution of the number of Episodes of ASBO by years

Year	Number of episodes of ASBO
2012	7
2013	4
2014	11
2015	4
2016	13
2017	12
2018	10
Total	61

### 2. Gender

Out of a total of 57 patients included in our study, 46 (80.7%) were males, and 11 (19.3%) were females. the sex ratio was 4.18



**Figure 01:** Gender distribution of the studied group

### 3. Age :

The mean age of our patients was 7 years and 5 months . and the age extremes ranged between 8 days and 15 years.

**Table II: distribution of patients by age groups**

Age group	Number (N)	%
Newborn (0 – 1 month)	1	1.75%
infant+ toddler ( > 1 month – 3 years )	9	15.79%
preschooler child ( > 3 – 6 years )	12	21.05%
school age child ( > 6 – 12 years )	23	47.37%
Adolescent ( > 12 years )	6	14.04%

## II. PAST MEDICAL HISTORY

### 1. Medical history

Only one patient of our studied group had a medical condition, he was treated for abdominal tuberculosis, and he had a peritoneal biopsy.


### 2. Surgical history

3 out of our 57 patients had neither medical nor surgical history. The remaining 54 patients were operated on for different surgical conditions.

36 of the 54 patients were operated on for appendicitis (63.16%). their surgical records have shown, that 27 patients have had perforated appendicitis (47.37%) and 9 patients with no perforation (15.79%). Four patients were operated on for Intussusception (7.02%). 3 had perforated bowel after abdominal trauma (7.02%). 3 patients had midgut volvulus (5.26%) . two patients had small bowel atresia (3.51%) . two others had omphalocele (3.51%). Diaphragmatic hernia (1.75%), sigmoid volvulus (1.75%) and peritoneal biopsy (1.75%) were found one time for each one. (the results are shown in the figure 03 bellow).

In our studied group, all patients who had surgical history have been operated on by laparotomy.

53 of 54 surgical procedures were performed in infracolic compartment while only one was performed in supracolic compartment for diaphragmatic hernia cure



**Figure02:** distribution of the previous surgical conditions leading to ASBO


### III. History of presenting illness

#### 1. Symptoms :

**Table III: distribution of ASBO symptoms in our patients**

Symptoms	Number of ASBO episodes	Percentage
➤ <b>Abdominal pain</b>	<b>61</b>	<b>100.00%</b>
• diffused pain	56	92 %
• Umbilical	5	7 %
➤ <b>Vomiting</b>	<b>61</b>	<b>100.00%</b>
• Bilious vomiting	37	61 %
• Nonbilious vomiting	24	39 %
➤ <b>Obstipation</b>	<b>59</b>	<b>97 %</b>


#### 2. the onset of symptoms :



**Figure 03: Distribution of the onset of symptoms**

### 3. Patient delay:

it ranged from 6 hours to 7 days, and the mean delay was 54 hours.



**Figure04:** Distribution of the time interval between the symptom's beginning and doctor visit

### 4. the Time period between the previous surgery and development of ASBO symptoms


The mean time period for the development of ASBO symptoms was 1 year. the extremes ranged between 7 days and 10 years

**Table IV:** Distribution of the time period between the surgery and the development of symptoms

Time period	Number of patients	percentage
[0 – 6 weeks]	13	24.07%
]6 weeks – 3 months]	16	29.63%
]3 months – 12 months ]	16	29.63%
beyond 1 year	9	16.67%

## IV. PHYSICAL EXAMINATION

### 1. General examination



**Figure 05:** Distribution of physical signs found in the general examination

### 2. Abdominal examination :

**Table V:** Distribution of the abdominal examination findings


Physical signs	Number of ASBO episodes	Percentage
➤ Abdominal distension	35	57 %
➤ Abdominal tenderness	61	100%
• Diffused	- 49	- 80 %
• Umbilical	- 9	- 15 %
• Right Iliac region	- 3	- 5 %
➤ Abdominal guarding	3	5%

## V. PARACLINICAL INVESTIGATIONS

### 1. Radiological investigations

#### 1.1. Abdominal X-ray (AXR)

The AXR has been performed in all patients of our studied group, and it showed distended loops of bowels and air–fluid levels (AFL) mostly central.



**Figure 06:** Distribution of the Air–fluid levels types found in our studied group



**Figure 07:** AXR of a 13 years old child showing central Air–fluid levels and dilated loops of bowel



**Figure 08:** AXR of a 7 years old showing peripheral Air-fluid levels and DLB



**Figure 09:** AXR of 12 years old child showing both central and peripheral Air-fluid levels

### **1.2. Abdominal ultrasonography**


The abdominal ultrasonography was performed in 9 patients. It showed dilated loops of bowel in all of them, and peritoneal effusion in 6 patients.

## **2. Biological tests**

9 patients have Hyperleucocytosis, one patient had severe anemia that needed a blood transfusion, and 14 patients have electrolytes imbalances including hyponatremia, hypochloremia and hypokalemia.

## VI. MANAGEMENT


out of 61 ASBO cases, 11 cases (18%) underwent immediate surgery without any previous conservative therapy trial . 50 cases ( 82 % ) were treated conservatively, and out of those patients, 16 cases (26% ) underwent delayed surgical exploration after unsuccessful conservative management.



**Figure 10:** distribution of the types of management in our studied group

### 1. Conservative management

In all the 50 episodes where the patients have been managed conservatively, the patients were treated according to the standard decompressive therapy which included nasogastric decompression, exclusion of oral intake, intravenous fluid, correction of electrolytes imbalance, analgesics, monitoring and frequent reassessment. 33 patients among them received water-soluble contrast agent (Gastrographin ) just after their admission along with the decompressive therapy, without waiting for a 48 hours delay.



**Figure 11:** distribution of conservative management protocols performed in our patients

1.1. Duration of the conservative management trial

**Table VI:** minimum, maximum, and the mean duration of the conservative management trial

Treatment	Duration		
	min	max	mean
conservative management Trial	1 day	8 days	2.7 days
• Standard decompressive therapy	1 day	7 days	2.1 days
• Gastrographin protocol	1 day	8 days	3 days


1.2. The outcome of the conservative management

**Table VII:** Distribution of conservative management outcomes

Treatment	Number of episodes		success rate
	Successful outcome	Unsuccessful outcome	
conservative management trial	34/50	16/50	68 %
• Standard decompressive therapy	13 /17	4/17	76 %
• Gastrographin prtocol	21 / 33	12 /33	64 %



**Figure 12:**Abdominal Radiographs of a 13 years old male patient. A- AXR obtained 3 hours after Gastrographin administration showing contrast in the stomach. B- AXR obtained 12 hours after Gastrographin administration showing the presence of the contrast in the large bowel




**Figure 13 :**Abdominal Radiographs of a 9 years old male patient , A- AXR showing Air-fluid levels and dilated loops of bowel . B- AXR obtained 12 hours after Gastrographin administration. C- AXR Obtained 18 hours after Gastrographin administration showing the contrast in the large bowel and the rectum .

## 2. Operative management

### 2.1. Timing

27 patients (44 %) out of 61 cases have been managed operatively, eleven (18 %) of them have had immediate surgical exploration after their admission, the remaining 16 patients (26 %) underwent delayed surgery after unsuccessful conservative management trial.



**Figure 14:** distribution of the operative management timing

### 2.2. Surgical approach

All operative managements in our studied group have been performed using Laparotomy

### 2.3. Surgical exploration

**Table VIII:** distribution of the surgical exploration findings

Surgical exploration findings	Number of patients	Percentage
Single bowel band	15	56%
Multiple bowel bands	12	44%
Adhesions	17	63%
Volvulus	6	24%
Gangrenous bowel	2	8%

## 2.4. Surgical procedure

**Table IX:** Distribution of the surgical procedures performed in our studied group

Surgical procedure	Number of patients	Percentage
<b>Adhesiolysis</b>	25	93%
<b>Retrograde decompression</b> (Milking of the intestinal content )	25	93%
<b>Intestinal resection and anastomosis</b>	2	7%
<b>Perforation repair</b>	0	0%
<b>Decompressive enterotomy</b>	0	0%

The intestinal resection was performed in two patients . for the first patient, it took away 25 cm of the ileum length and left out 7 cm of terminal ileum downstream . for the second, the resection took away 15 cm of the ileum length and left away the last 10 cm of the terminal ileum.

## 2.5. Post-operative outcomes

One patient had postoperative sepsis that needed a transfert to critical care departement .

# VII. Evolution :

## 1. Length of hospital stay :

**Table X:** Distribution of the length of hospital stay by therapeutic groups

Group	length of hospital stay		
	min	max	mean
➤ <b>Conservatively managed group</b>	1 day	8 days	3 days
• <b>Standard decompressive therapy</b>	1 day	7 days	2.1 days
• <b>Gatrogaphin protocol</b>	1 day	8 days	3.5 days
➤ <b>Operatively managed group</b>	4 days	13 days	6.1 days
➤ <b>All the studied group</b>	1 day	13 days	4.4 days

## 2. Recurrence


There have been 4 cases (6.5 %) of ASBO recurrence, 3 of them were previously managed conservatively ( 2 patients were managed with standard decompressive therapy and one received WSCA), the remaining case was managed operatively. During the recurrent episodes, the 3 patients who have been previously managed non operatively underwent surgery after unsuccessful conservative trial, while the patient who had been previously managed operatively was successfully managed non operatively.

**Table XI : Features of the recurrent cases of ASBO**

Cases	Case 1	Case 2	Case 3	Case4
<b>Gender</b>	Male	Male	Male	Male
<b>Age</b>	11 years	12 years	1.5 year	7 years
<b>The Previous surgical condition</b>	Perforated Appendicitis	Perforated Appendicitis	Intussusception	Intussusception
<b>Management of the first episode of ASBO</b>	Immediate surgery	Standard conservative management	Standard conservative management	Conservative management using Gastrographin
<b>Time of recurrence</b>	4 months	6 months	3 days	6 days
<b>Management of the recurrent episode of ASBO</b>	Standard conservative management	Delayed surgery (after failure of conservative management)	Delayed surgery (after failure of conservative management)	Delayed surgery (after failure of conservative management)

## 3. Mortality

Two cases of death (3.5%) were registered in our studied group. One of them died after 24 hours of conservative management, from a sudden cardiopulmonary arrest secondary to massive inhalation of gastric content. The second one was managed operatively after 72 hours of unsuccessful conservative management; he died in the critical care unit from postoperative sepsis.



# *DISCUSSION*



## I. Epidemiological data

### 1. GENDER

**Table XII:** the sex ratio reported by authors

Author	number of cases with ASBO	sex ratio (M/F)
J. Janik & al ; Canada ; 1981 [13]	131	1,2
Lautz TB & al ; USA 2011 [14]	2089	1,7
Al-Salem ; Saudi Arabia 2012 [15]	44	1,7
A. Eeson & al ; Canada 2010 [16]	151	2,2
Nasir & al ; Nigeria 2013 [17]	618	1,8
Akgur & al ; Turkey 2017 [18]	181	2
CY Lee & al ; Taiwan 2014 [19]	33	1.4
F. Linden; USA 2019 [20]	12	5
Z. Bouhnoun ; Morocco (Fes) 2016 [21]	43	1.2
our study	61	4.1

Studies conducted in adults, noted that gender was not a risk factor of ASBO . some studies reported males predominance [22–24], while others recorded predominance of females [25,26]

in pediatric patients, all the reviewed studies reported a predominance of males with various sex-ratio.

## 2. AGE

**Table XIII : the mean age and age extremes reported by authors**

Author	Number of cases with ASBO	mean age	age extremes
A. Eeson & al ; Canada 2010 [16]	151	6.6 years	2 weeks – 17 years
Nasir & al ; Nigeria 2013 [17]	618	4.5 years	3 months – 15 years
Bonnard & al ; France 2010 [27]	8	7 years	4.4 years – 13.1 years
Al-Salem ; Saudi Arabia 2012 [15]	44	5.4 years	1 month – 12 years
C Festen; Netherland 1982 [7]	31	5,2 years	3 weeks – 16 years
LB Chirdan ; Nigeria 2011 [29]	22	7 years	21 days – 14 years
N. Erturk& al ; Turkey 2017 [28]	45	7.75 years	6 days – 14 years
Lautz TB & al ; USA 2011 [14]	2089	12,6 ans	6,8 years – 18,4 years
J. Janik & al ; Canada ; 1981 [13]	131	6 years	1 month – 17,9 years
Z. Bouhnoun; Morocco Fes 2016 [21]	43	9.5 years	9 months – 15 years
our study	61	7.5 years	8 days – 15 years

Adhesive small bowel obstruction can occur at any age [30]. Young et al.[31] reported a higher incidence of ASBO in infants under 1 year of age, irrespective of the initial site or indication for laparotomy. Conversely, the study conducted by N. Erturk et al [28] showed that 82.5% of ASBO occurred among children older than one year, while only 17.7% occurred in infant under 1 year . in the other hand Grant et al [6] found that there is no difference in readmission rates for ASBO between younger and older children when comparing the organ on which surgery was initially performed.

In our study, the mean age of the children with ASBO was 7.5 years, though some studies reported a mean age older or younger than our patients. Nasir et al [17] reported a mean age of 4.5 years, and in the series of Lautz [14] the mean age was 12.6 years.

### **3. PAST MEDICAL HISTORY**

#### **3.1. Medical history**

Fewer are the studies that reported the existence of a previous medical condition in their patients with ASBO. XY Cao [32] Reported 51 cases of ASBO in patients that were being treated for abdominal tuberculosis, and he also reported that Abdominal tuberculosis is an important and increasingly common cause of acute bowel obstruction.

In our study, only one child among our young patients had a previous medical condition; he was treated for abdominal tuberculosis. The same patient also had a peritoneal biopsy two months before the beginning of ASBO symptoms, which would most likely be the cause of those adhesions.

### 3.2. Surgical history

**Table XIV:** the mean previous surgical conditions leading to ASBO in children reported by authors

Author	Previous surgical conditions
<b>Akgur&amp; al ; turkey 1991 [18]</b>	1/ Appendicitis = 28.73% 2/ Abdominal trauma = 19.3 % 3/intussusception = 11.6%
<b>Feigin&amp; al ; 2010 [33]</b>	1/Appendicitis= 15.5 % 2/ Congenital bowel defect = 13.8% 3/ Abdominal tumor = 13.2 %
<b>Nassir &amp; al ; Nigeria 2013 [17]</b>	1/ typhoid intestinal perforation= 24.1% 2/Intussusception= 20.7 % 3/ volvulus and malrotation = 17.2%
<b>N. Erturk&amp; al ; Turkey 2017 [28]</b>	1/ Appendicitis *(without perforation) = 15.5% *(with perforation ) = 20% 2/ megacolon + ARM = 15.5% 3/ small bowel atresia = 11%
<b>Young &amp; al ; USA 2007 [31]</b>	1/ Appendicitis ( without perforation )= 3% *(with perforation ) = 13% 2/ Malrotation = 8% 3/ Hirschsprung disease = 6.5%
<b>C.-Y. Lee &amp; al ; Taiwan 2014 [19]</b>	1/ Appendicitis = 37.5 % 2/ Hirschsprung disease = 12.5% 3/ Intussusception= 12.5%
<b>Z. BOUHNOUN ; Morocco (fes) 2016 [21]</b>	1/ Appendicitis * (without perforation ) = 11.5% *( with perforation ) = 60.5 % 2/ Intussusception = 13 % 3/ Volvulus and malrotation = 5%
<b>Our study</b>	1/ Appendicitis *(without perforation)= 15.79 % *(with perforation ) =47.37 % 2/ Intussusception = 7 % 3 / abdominal trauma = 7% 4/ volvulus and malrotation = 5%

Various previous surgical conditions have been reported in children with ASBO. Vijay [34] found that Hirshprung disease and Intussusception were associated with high rates of ASBO . in Nasir’s Study[17] typhoid intestinal perforation was the frequent previous condition.

Appendicitis was reported by the majority of authors [18,33,28,19,21] as the commonest previous condition associated with ASBO in children. Moreover the incidence of ASBO was higher after perforated appendicitis than non perforated one . some studies [13, 31] conducted in children to assess the risk after appendectomy , they noted a low incidence of ASBO after non perforated appendicitis ,elseways the incidence after perforated appendicitis was relatively higher , Young & al [31]reported a rate of 0.3% after non perforated appendicitis versus 3% after perforated appendicitis . this can be explained by the high level of inflammation that occurs during perforated appendicitis and the large laparotomy incision that is performed in this case, which raises the risk of adhesions formation [1]

In our study, the majority of the children with ASBO have had appendicitis.47.37% of cases have had perforated appendicitis and 15.79 % without perforation.

Many studies conducted in African countries [17,21, 34] reported Intussusception as the second previous condition in children with ASBO. Nasir & al [17]reported that 20.6% of the children with ASBO have had previous Intussusception . in our study intussusception was noted in 7 % of the children.

Grant et al [6] found that operation on the ileum of children is followed by a high risk of readmission for adhesionrelated problems. This risk is much higherand it can reach up to 25% when Ileostomy formation and closure is performed, that goes further to confirm our findings of high incidence of ASBO in children with previous midgut malrotation or small bowel perforation resulting of abdominal trauma, their incidence in our study were respectively 5% and 7% . those findings are consistent with other studies reports[17,18,21,31]

Aguayo & al [4] conducted a study on the risk of bowel obstruction after treatment of abdominal tumours; he concluded that Bowel obstruction is relatively uncommon after intra-abdominal malignancies in children, with an overall incidence of 3.7 %. However, Wilms ' tumour, rhabdomyosarcoma and Burkitt' s lymphoma appear to be associated with the highest risk of bowel obstruction in comparison with other intra-abdominal malignancies.

Tree children (5%) enrolled in our study have no medical nor surgical history, and though they presented ASBO . all of them underwent surgical exploration which has confirmed the presence of congenital adhesions. The incidence of adhesions without previous surgery has been reported to range between 3% and 28 % as determined by autopsy [36,37,38]. Congenital adhesion band is a very rare condition[38] which make a few existing studies that reviewed small bowel obstruction by Congenital adhesions. KH Yang [38] reported that 5.9 % of the patients with ASBO included in his research had congenital adhesions, including 2/3 of them that were children. And unlike adult patients, pediatric patients with congenital ASBO showed a high proportion of early operation and bowel resection [38].

### **3.3. The time interval between the previous surgery and the development of ASBO**

Obstruction can occur anytime from the early post-operative period to many decades later after the surgical procedure [39], a Swedish study conducted by RE. Anderson in adult patients showed that ASBO can occur even 30 years after laparotomy[40]. In pediatric patients, most of ASBO occur within the first year after surgery [31,34,41,42]. Akgur & al [18] reported that most of the ASBO occurred less than 3 months after laparotomy, and those episodes were associated with a high success rate of conservative management which was 81% versus 59 % when the time elapsed after the laparotomy is more than 18 months.

In our study, 53.7% of ASBO episodes occurred in less than three months after the surgical procedures, and about 80.3% occurred in less than one year, which is consistent with the previous findings in the literature.

## II. PATHOPHYSIOLOGY

### 1. Etiopathogenesis of adhesions

#### 1.1. Post-operative adhesions formation [43,35,45,48]

Peritoneal adhesions may be acquired or congenital; however, most are acquired as a result of peritoneal injury, which occurs by inflammatory conditions, intraperitoneal infection, abdominal trauma, or most commonly abdominopelvic surgery.

The mechanisms of post-operative adhesion-genesis are common to all acquired peritoneal adhesions. They involve mesothelial surface disruption with subsequent fibrino-coagulative and inflammatory signalling processes.

When peritoneal serous is injured, the process of repair or healing occurs in two ways; the first one is the usual way which leads to the resolution of the injury and peritoneal repair. The second way is pathological and it leads to persistent adhesions formation.

Immediately after injury, there is bleeding and an increase in vascular permeability with fluid leakage from injured surfaces. Simultaneously, a posttraumatic inflammatory response occurs, with infiltration of inflammatory cells, the release of pro-inflammatory cytokines and activation of the complement and coagulation cascades. The activation of the coagulation cascade results in the formation of thrombin, which is necessary for the conversion of fibrinogen to fibrin, which functions to restore injured tissues and, once generated, is deposited along peritoneal surfaces. Fibrin is a tacky substance and causes adjacent organs or injured serosal surfaces to coalesce.

Under normal circumstances, the formation of a fibrin matrix during wound healing is only temporary and degradation of these filmy fibrinous adhesions by locally released proteases of the fibrinolytic system occurs within 72 hours of injury. Fibrinolysis allows mesothelial cells to proliferate and the peritoneal defect to be restored within 4 to 5 days, preventing the permanent attachment of adjacent surfaces. This process has the key role in tissue remodelling and repair.

If fibrinolysis does not occur within 5 to 7 days of peritoneal injury, or if the local fibrinolytic activity is reduced, the fibrin matrix persists. If this occurs, the temporary fibrin matrix gradually becomes more organized as collagen-secreting fibroblasts and other reparative cells infiltrate the matrix. The organization of fibrin bands over time and their transformation into mature fibrous adhesions is what enables them to persist. These "mature" adhesions are not simply composed of connective tissue; studies have demonstrated that, over time, they become highly organized cellular structures that contain arterioles, venules, capillaries and nerve fibres in addition to collagen.




Figure 15: Illustration image summarizes the process of adhesion formation [108]




FIG. 1. Biological events involved in peritoneal tissue repair and adhesion formation. PAI-1 = plasminogen activator inhibitors group 1; tPA = tissue plasminogen activator; uPA= urokinase-like plasminogen activator.

Figure 16: Biological events involved in peritoneal tissue repair and adhesion formation[43]

### **1.2. Congenital Adhesions formation**[38,47]

The congenital adhesions and bands have no relation to an intra-abdominal process (previous laparotomy, inflammatory diseases, peritonitis, embryogenic remnants, etc.), they are considered to have embryologic bases, such as incomplete regression of the fetal vitelline circulation, or a genetic defect that impairs embryogenesis. Furthermore, congenital adhesions might also be a result of intrauterine exposure to certain infectious agents or ischemic events.

## **2. Mechanisms of obstruction** [ 44,45,46 ]

The ASBO is a mechanical obstruction, the mechanism of this obstruction can occur either by direct extrinsic compression . or by volvulus, during which, the loop of intestine twists around itself and the mesentery that supports it, resulting in bowel obstruction and later on ischemia and gangrene due to arterial occlusion

## **3. Consequences of intestinal obstruction** [ 43,8,47,48 ]


The pathophysiologic alterations produced by the intestinal obstruction are both local and general. With mechanical occlusion of the lumen, the intestine above becomes distended, resulting from an accumulation of gas and fluid within its lumen. This distention is enhanced by swallowed air, diffusion of gas via the bloodstream, multiplication of bacteria within the stagnant intestinal contents, and hypersecretion of the injured gut.

When intraluminal pressure exceeds capillary and venous pressure in the bowel wall, the intestinal absorption and lymphatic drainage decrease, the bowel becomes ischemic when capillary blood flow stops, allowing bacteria to pass into the peritoneum; from that point, it passes into the bloodstream, leading to septicemia by a process known as bacterial translocation. The peritoneal fluid is continuously secreted by the visceral peritoneum and absorbed by the parietal peritoneum, mostly in the diaphragmatic abdominal surface, where the diaphragmatic pores can be distended as much as 3 times their normal size to allow the passage of bacteria. The colonized fluid is then transported via the lymphatic channels into the thoracic duct, which drains in the jugulosubclavian angle, allowing bacteria to enter the circulation and causing septicemia.

Perforation can develop as the ischemia leads to bowel necrosis. First, lymphatic obstruction occurs because of the lesser pressure in these vessels. This is followed by a venous obstruction, which accelerates the edema process because blood enters the affected bowel segment but does not have a drainage route. Finally, the continuous increase in the bowel wall pressure blocks the arterial vessels, leading to ischemic necrosis and perforation.

The massive third spacing of fluids results from peritoneal reflex hypersecretion. Rapidly leads to shock, contributing to morbidity and mortality. This sequence may occur more quickly in a closed-loop obstruction with no proximal escape for bowel contents.

In addition to the local effects produced by obstruction, most profound systemic alterations occur. These may be summarized as a rapidly occurring dehydration with oliguria, hemoconcentration, hypochloremia, hypopotassemia, alkalosis, retention of certain nitrogenous substances, and alterations of the serum proteins.



Pathophysiologic changes of small-bowel obstruction.

**Figure 17:** illustration of pathophysiological changes of small bowel obstruction [49]




Figure 18: consequences of small bowel obstruction

### **III. CLINICAL FEATURES [8,44,45,46]**

#### **1. SYMPTOMS**

##### **1.1. Abdominal pain**

Pain is the earliest symptom to appear; it can be acute or progressive . and most commonly of a colicky character. Furthermore, in children, the pain is associated With motor agitation and crying, and as the obstruction progress, The child becomes increasingly lethargic.

Neither the site nor the type of pain can determine the exact location of the obstruction.

##### **1.2. Vomiting**

Vomiting is as early as the site of obstruction is more proximal; initially, the emesis may be nonbilious, but with time it progresses to bilious or event feculent emesis . in 30% of cases the vomiting is associated with relief of abdominal pain.

##### **1.3. Obstipation**

With partial obstruction, there still be a continuous passage of flatus and small stool. But in children with complete obstruction both of flatus and stool ceases. Interestingly, in early stages, diarrhea may be present and might confuse the clinical picture.

In our study, abdominal pain and vomiting were present in all children (100% ), and the obstipation was found in 59/61 patients (97%)

## 2. Physical signs

### 2.1. Systemic signs

In the early stage, children with SBO could present Tachycardia along with palor and motor agitation, which can be related to high pain intensity. Later on, when the patients are seen in a late-stage, they can present tachycardia, Fever, hypotension, and dehydration, those signs indicate commonly circulatory collapse and bowel gangrene.

In our study fever was found in 5/57 patient ( 8%), dehydration in 12/57 patients (24.5%) and the tachycardia in 5 patients (8%)

### 2.2. Abdominal examination

#### ➤ **Inspection**

The inspection of the abdomen notices the amount of distension which is greater in case of lower small bowel or colonic obstruction. While in proximal bowel obstruction it might not be so marked .furthermore visible peristalsis under the abdominal wall can be seen . and also abdominal scars related to previous surgery but in case of congenital ASBO abdominal scar is missing.

#### ➤ **Palpation**

Palpation usually finds abdominal tenderness, diffused or localized . Less commonly, abdominal guarding can be found, which highly refers to bowel necrosis or event peritonitis in case of diffused abdominal guarding.

#### ➤ **Percussion**

Is of a little value except for tympanic resonance that confirms abdominal distension or dullness over the flanks which suggestive of peritoneal effusion.

➤ **Auscultation**

Also, it is of a very little help. Initially, it noticed hyperactive bowel sounds due to hyperperistalsis then decrease.

**2.3. Rectal examination**

A rectal exam must be performed. This gives important information, including the absence of stool, and possibly blood.

## **IV. PARACLINICAL FEATURES**

### **1. Radiological investigations**

Radiologic imaging plays a crucial role in the diagnosis and management of patients with suspected SBO. A myriad of radiologic investigations are available, and the evaluating physician must choose in a reliable, cost-effective way. [46]

#### **1.1. Abdominal X-ray [44, 50,51,52,53]**

After a past medical history and physical exam, the initial radiographic study would be plain abdominal radiographs in orthostatic position .even though an upright film is optimal, lateral decubitus or cross-table lateral films could be acceptable substitutes in bedridden patients.

In ASBO, the abdominal radiographs can classically show the following features.:

- dilated loops of small bowel proximal to the obstruction >2.5–3 cm
- predominantly central dilated loops
- valvulae conniventes are visible
- air–fluid levels in erect view study, especially suspicious if >2.5 cm in length and in the same loop of bowel but at different heights
- free intraperitoneal air can be found in case of bowel perforation

Obviously, any or all of these findings may be present, but the skilled clinician should be able to correlate these radiograph findings with the clinical picture. A recent study evaluating plain radiographs in patients with suspected SBO demonstrated good accuracy, with a mean sensitivity of 83% for correctly identifying a SBO[51]. while it has a low specificity in diagnosing an SBO since both mechanical obstruction and functional bowel disorders may appear identical [46]

## 1.2. Abdominal Ultrasonography [52,54,55]

Ultrasound is a Bedside test also an alternate modality, but really of limited use, especially when intestinal dilation is present and air-fluid levels limit the effectiveness of the study. Adhesions and band are not seen in ultrasound images, but it shows many suggestive findings of small bowel obstruction :


- dilated bowel loop (diameter >2.5 cm)
- ineffective peristalsis (results in "whirling" appearance of intra-luminal contents)
- the prominence of the valvulae conniventes (present in dilated jejunal loops)

Furthermore, ultrasound can help to find suggestive signs of bowel ischemia/ infarction that will need urgent surgical evaluation :

- extra-luminal free fluid ( the "pointy" triangular appearance of inter-loop free fluid is often referred to as the tanga sign )
- loss of peristalsis
- bowel wall thickening >3 mm (with effacement of mural architecture)
- intramural bowel gas



**Figure 19:** abdominal ultrasound image showing, dilated small bowel loop, with diameter 3.24 cm (> 2.5 cm ) [104]



**Figure 20:** Ultrasound image showing a dilated loop of bowel and extra-luminal free fluid known as "Tanga sign " [56]


### 1.3. Abdominal CT scan [52,57]

Abdominal CT scan, especially with administration of oral or intravenous contrast, performs better than plain Xray in finding the transition point, evaluating the severity of obstruction, identifying the cause of obstruction, and recognizing complications such as ischemia, necrosis, and perforation [58]. The sensitivity, specificity, and accuracy of CT scans for ASBO diagnosis are very high; they are, respectively, from 90% to 94%, 96%, and 95% [59]. The ASBO Features on CT may include:

- dilated small bowel loops >2.5–3 cm
- normal caliber or collapsed loops distally
- small bowel feces sign which can be defined by the presence of feculent matter mingled with gas bubbles in the lumen of dilated loops




**Figure 21: Axial abdominal CT showing the transition point between dilated bowel loop and collapsed loop (red arrows) [61]**



**Figure 22:** Axial abdominal CT showing small bowel feces sign (blue arrow) [62]

CT scan can also show characteristic features of Closed-loop obstruction ( when a bowel loop of variable length is occluded at two adjacent points along its course) :

- radially distributed, C or U-shaped small bowel loops
- " Fat notch sign":reflects insinuation of mesenteric fat at an area of adhesions with focal caliber change
- "beak sign" corresponding abrupt luminal transition and luminal constriction
- two adjacent collapsed loops of bowel



**Figure 23:** Axial abdominal CT of an adult patient showing closed-loop obstruction caused by adhesive bands[63]




**Figure 24:**coronal reconstruction of an abdominal CT scan showing the "Fat notch sign" ( white arrow) [64]




**Figure 25** : coronal reconstruction of an abdominal CT scan showing the “beak sign” (double arrow)[64]

When the diagnosis of closed-loop obstruction is delayed, it may be associated with intestinal ischemia which is known as bowel strangulation. It is considered as a dangerous complication since it is associated with high mortality. Strangulation presents with non-specific features ,it include[60]:

- thickened and increased attenuation of the bowel wall
- halo or target sign
- pneumatosis intestinalis: gas in the intestinal wall
- pneumatosis portalis: gas in the portal vein or in the mesenteric vein
- localized fluid or hemorrhage in the mesentery
- variable amounts of peritoneal fluid



**Figure 26:** Axial Abdominal CT showing the difference between normal bowel wall enhancement and ischemic bowel wall enhancement [62]



**Figure 27:** Axial abdominal CT showing Intramural bowel gas or pneumatosis intestinalis[65]

However, abdominal CT scan should not be routinely implemented in the diagnosis making process except when clinical history, physical examination, and plain film were not convincing for ASBO diagnosis, since these are readily available, less expensive, expose the patient to less radiation, and may highlight the need for abdominal CT in some patients . [66]

## 2. Biological tests

The minimum of laboratory tests includes; blood cell count, lactate, electrolytes, CRP, and BUN/creatinine. Currently, available laboratory tests do not contribute to the diagnosis of SBO, but they can confirm the clinical suspicion of volume depletion, guide volume resuscitation, and assist recognition of bowel ischemia. [67]

Elevated Leukocytosis count and CRP has been implicated as a marker of strangulation, although sensitivity and specificity of these tests are relatively low [68]

Electrolytes are often disturbed in patients with a bowel obstruction; in particular, hypochloremia, hypokalemia, and metabolic alkalosis are frequently found. Dehydration and Hypovolemia may manifest as azotemia with an elevated blood urea nitrogen (BUN) to creatinine ratio (BUN/creatinine), and it could result in acute kidney injury [71]

L-lactate, lactate dehydrogenase (LDH) and creatine kinase (CK) can rise due to hypoperfusion of the intestinal tissue. L-lactate level increases sensitivity for detecting bowel ischemia and is considered a reliable indicator for emergency surgical intervention. In contrast, LDH and CK raised levels are therefore unspecific [69,70]

## **VI. MANAGEMENT**

### **1. The Goals of management [44,46,47,71]**

The management of Adhesive small bowel obstruction in children consists of the following goals :

- ✓ Resuscitation, restoring of normal fluid volume and the correction of electrolytes imbalances
- ✓ Relief of pain and discomfort
- ✓ Achieving resolution of the obstruction and minimize morbidity and mortality

### **2. Therapeutic approaches and Indications**

#### **2.1. Conservative management**

##### **2.1.1. Standard conservative management without WSCA**

The cornerstone of non-operative management is gastro-intestinal decompression with a nasogastric tube and bowel rest by nil peros. Moreover it consists of, intravenous fluid hydration, correction of electrolytes imbalances, nutritional support, Strict input and output measurements, symptomatic treatment, and Prophylactic antibiotics that should be administered as the clinical situation dictates .in addition to that patients should be Frequently reassessed to make sure that the patient is not developing suggestive signs of complications . **[71,72]**

**Table XV: Distribution of ASBO's therapeutic approaches and the success rate of conservative management according to studies reports**

Author	number of patients with ASBO	number of ASBO episodes	number of episodes underwent immediate surgery	Number of episodes managed conservatively		the success rate of conservative management	
Akgür (1991) [18]	181	230	81 (35 %)	149 (65 %)		73,8 % (110)	
Vijay (2005) [34]	69	74	5 (7 %)	69 (93 %)		52,2 % (36)	
Eeson (2010) [16]	151	165	32 (19 %)	133 (81 %)		16 % (26)	
Osifo (2010) [73]	21	21	0 (0%)	21 (100%)		0% (0)	
Feigin (2010) [33]	128	174	0 (0%)	174 (100%)		63% (109)	
Bonnard (2010) [27]	24	24	0 (0%)	24 (100 %)		58 %	
				* without WSCA 16	* with WSCA 8	* without WSCA 50 % (8)	*with WSCA 75 % (6)
Nasir (2013) [17]	29	34	0 (0%)	34 (100%)		47% (16)	
Bouhnoun (2016) [21]	43	49	1 (2 %)	48 (98 %)		18 % (9)	
our study	57	61	11 (18 %)	50 (82 %)		68% (34)	
				* without WSCA 17 (28%)	* with WSCA 33 (54%)	* Without WSCA 76 % (13)	*With WSCA 64 % (21)

The paradigm for the initial management of children with ASBO is similar to that for adults and has not changed appreciably in the last two decades [39]. The last update of guideline that was written by Broek et al under the auspices of the World Society of Emergency Surgery recommends that Non-operative management should always be tried in patients with adhesive small bowel obstruction, Unless there are signs of peritonitis, strangulation, or bowel ischemia [72]

Akgur et al [18] suggested the use of a conservative trial for managing the patients in the absence of fever, leukocytosis, localized tenderness, and complete obstruction. And Stewardson [74] suggested immediate surgery for patients with any of 4 those classic predictors (fever, tachycardia, leukocytosis, localized tenderness).

Inconsistently Eason reported in his study that 46% of patients who resolved without surgery, had at least one of the classic predictors at presentation. According to him, those clinical criteria promote an overly aggressive surgical management, and only fever and leukocytosis could predict the need for operative management. [16]

There is no evidence for the optimal duration of non-operative treatment, but most authors consider a 72-h period as safe and appropriate [72,75,76,77,78]. Conservative treatment can be continued for more than 72 h in cases with persistent high output from a decompression tube, but no other signs of clinical deterioration. Lautz et al [39] recommended an observation period of 24 to 48 hours in stable infants/ young children less than 1-2 years , and 48 hours for children greater than 1-2 years of age . in the absence of clinical indicators suggesting improvement, observation periods greater than 48 hours carry a higher risk of bowel ischemia and necrosis, and the likelihood of resolution of the ASBO is extremely low.

The objective criteria that identify those patients who are likely to respond to conservative treatment does not exist. Eason [16] suggested that the absence of leukocytosis and older age are predictors of successful conservative management. Some authors[71,75]proposed the following as strong predictors of NOM failure:

- ✓ The presence of ascites
- ✓ complete ASBO (no evidence of air within the large bowel),
- ✓ increased serum creatine phosphokinase
- ✓  $\geq 500$  mL from the nasogastric tube on the third NOM day.

The success rate of the usual conservative management without WSCA ranged from 0% to 73.8 %. four of those studies included in our review [18,27,33,34] reported that conservative treatment was successful in more than 50% of cases . on the other hand , it was successful in none of the cases in the study reported by Osifo [73], however the author acknowledged that the study was conducted in a poor resource country that lacks adequate facilities and manpower which appears to have limited the capacity for comprehensive conservative treatment and monitoring . Despite the presence of resource, Eeson et al [16] reported a much lower rate of conservative treatment success (16%) than reported in the other studies. The reason for this much lower rate of treatment success is not apparent, but would suggest that children in that study may have had more severe ASBO episodes than those in the other studies. Further studies are needed to identify the characteristics of children with ASBO who are most likely to respond positively to conservative treatment.

In the present study, the success of the standard conservative management without WSCA was observed in 13 cases among 17 cases that were treated according to the standard method, with a success rate of 76 % . which is consistent with the published reports

#### **2.1.2. Conservative management with Water-soluble contrast agent**


Water-soluble contrast agent has a diagnostic and therapeutic role [71,79]. Gastrografin is the most commonly utilized contrast medium. It is a mixture of sodium diatrizoate and megluminediatrizoate. Its osmolality is 2150 mOsm/L. Which is higher than the osmolality of extracellular fluid. It promotes shifting of fluid into the bowel lumen and increases the pressure gradient across an obstructive site. The bowel content is diluted, and in the presence of the wetting agent, the passage of bowel contents through a narrowed lumen is facilitated [79]. Gastrografin also decreases edema of the small bowel wall, and it may also enhance smooth muscle contractile activity that can generate effective peristalsis and overcome the obstruction

[71]. Gastrografin is water-soluble and relatively safe even if the obstruction is complicated by perforation. Complications from the use of Gastrografin in small bowel obstruction are rare, although anaphylactic reactions and lethal aspiration have been described[80,81]

Barium has also been used to evaluate adhesive small bowel obstruction; it is not as easily diluted by the enteric fluid as Gastrografin and provides a better mucosal image on radiography. However, a barium study can be risky because it may become inspissated and completely obstruct the bowel. Barium may spread into the peritoneal cavity if perforation occurs, a condition that is potentially lethal[79]

Contrast agent administration should be preceded by infusion of intravenous Ringer solution or Isotonic saline which would be administered first as boluses based on urine output then as a maintenance infusion based on body weight[27] . to guard against dehydration caused by the effect of the contrast medium. The patients should be monitored, and their electrolytes imbalances corrected prior to using Gastrografin . after dilution with an equal volume of isotonic solution, The prepared mixture is given according to the age 30-50 mL for under 10 years, and the other patients receive 100 mL. Through the nasogastric tube and kept closed for 3 h and then reopened. [19,27]

Patients should then have serial plain orthostatic abdominal radiographs at 3, 6, 12, and 24 h after contrast administration. after Gastrografin administration. the Management is considered successful when the contrast is visualized in the cecum and a clinical improvement is noted. Once successful management is achieved, the nasogastric tube is removed and feeding is initiated. If the contrast agent failed to reach the cecum, complete obstruction is diagnosed and patients will be scheduled for surgical intervention[19,20,27]



**Figure 28:** A-Abdominal radiograph of one of our patients ,obtained 12 hours after Gastrographin administration showing dilated bowel loop and Air-fluid levels filled with Gastrographin. B- Abdominal radiograph obtained 24 hours after Gastrografin administration showing that the contrast had reached the large bowel (hausters).

Many studies have been performed to evaluate the role of water-soluble contrast in the management of ASBO in adult patients[82,83,84]. They concluded that the contrast agent has great diagnostic benefit in assessing the indication for surgical intervention and the timing of surgery. The use of Gastrografin in ASBO patients helps to promote an earlier resolution of obstruction and also reduces the length of hospital stay compared with standard conservative management. However its role in reducing the rate of laparotomies remain inconclusive . in the other side, other studies that were conducted by Assalia [85] and, Di Savario [86] they found that Gastrografin decreased the fraction requiring surgery. Branco et al [87] and Abbas et al [83] reported that the appearance of water-soluble contrast in the colon within 24 hours after administration could accurately predict the resolution of ASBO with a sensitivity and specificity that reach 96 %. Atahan et al. [88] concluded that conservative treatment was recommended for

patients in which contrast medium is observed in the right colon within 8 hours following administration, regardless of the presence of obstruction signs.

**Table XVI:** the success rate of WSCA protocol according to studies reports

Author	Number of ASBO cases	Number of cases managed conservatively using WSCA	the success rate of conservative management using WSCA
Bonnard (2010) [27]	8	8	6 / 8 (75%)
Chee-Yew Lee (2014) [19]	33	19	16 / 19 (84%)
Abdelkader (2011) [89]	35	12	8 / 12 (66.6 %)
AF. Linden (2019) [20]	12	12	10 / 12 (83 %)
<b>Our study</b>	<b>61</b>	<b>33</b>	<b>21 / 33 (64 %)</b>

Reviewing the literature. We found 4 studies which have been conducted to evaluate the Role of WSCA in the management of ASBO in young patients. In all those studies included in our review, the success rate of this approach was high, and it ranged from 66.6% to 84 %.

All authors [19,20,27,89] reported the valuable role of WSCA in ASBO management. Bonnard [27] and Linden [20] administered the water-soluble contrast early after admission, conversely to the other related studies which delayed the administration until the fail of the 48 hours trial decompressive therapy [19,89]. Those studies conducted by Bonnard and Lindon were the only studies which have directly compared standard GI decompression versus the use of water-soluble contrast in the pediatric population they reported that the rate of success was relatively high in the water-soluble group compared to those treated with standard decompressive therapy, their success rate were accordingly 75% and 83% for WSCA groups versus 50% and 55% for the usual decompressive therapy groups. Furthermore, the length of hospital stay was also shorter in the water-soluble groups.

In our study, the success rate in the group treated with WSCA was 64 %, which is consistent with the studies reviewed. However when compared to our standard decompressive therapy group, this rate was relatively low (64% versus 75%), this result can be related to many limitations including the small number of patients, the difference in numbers in the two compared groups, and the variety of the previous surgical conditions leading to ASBO. Moreover, there was no standard time for the administration of the water-soluble contrast in our studied patients and in some cases a delay of administration, which may have influenced the therapeutic effect.

## 2.2. Operative management

### 2.2.1. Timing of surgery

The operative management of ASBO can be performed early or delayed after unsuccessful conservative management. There is an old dogma “Never let the sunrise or set on a small bowel obstruction” was based on the concern that delay in operation was associated with an increased risk of bowel ischemia, the need for bowel resection, and subsequent complications.[39,]

Emergent operative management is indicated for patients presenting with signs of bowel ischemia, which may include localized tenderness on exam, tachycardia, fever, laboratory studies indicating leukocytosis [39,18,74]. Eason [16] reported that only Fever and Leukocytosis were the only reliables that predicted the need for immediate surgery. The rate of patients who underwent immediate surgery ranged between 5% and 35 % [16,18, 34,89 ]. in our study 18 % of the patients were operated on immediately after their admission which is in accordance with the literature .

On the other hand, delayed surgery is performed after the failure of the conservative management . in the reviewed studies the rates of patients who underwent delayed surgery after failure of non-operative management varies between 26% and 100% [18,33,73]. In our studied group, 32% of the patients underwent delayed surgery, similarly to the rate reported by Fegin [33] which was 33 %.

Some authors indicate that there is a correlation between the timing of surgery and the risk of bowel ischemia. Feigin [33] reported that 31% of the patients managed operatively underwent bowel resection, and none of the patients who were operated on before the 16<sup>th</sup> hours needed bowel resection, he also reported that the rate of bowel resection in patients who were operated on in the first 48 hours was 12 %, which was lower compared with a rate of 27 % of bowel resection encountered in patients operated on lately after the 48 hours. In light of those findings, Feigin [33] concluded that the index of suspicion for compromised bowel rises after 16 hours, and the decision for surgery should be made at approximately 48 hours; moreover the observation should probably not be prolonged more than 72 hours.

### **2.2.2. Surgical approach**

Abdominal exploration through laparotomy has been the standard treatment for adhesive small bowel obstruction. In recent years, however, laparoscopic surgery for ASBO has been introduced [72]. The potential benefits of laparoscopy include less extensive adhesion reformation, earlier return of bowel movements, reduced post-operative pain, and shorter length of stay [90,91,92]


Aguayo et al [4] reviewed an experience with the laparoscopic approach for SBO in 34 children. They found that the laparoscopic approach was safe, but that 32% required conversion to a laparotomy, which can be due to an inability to find transition point, inability to achieve pneumoperitoneum, dense adhesions and gangrenous intestine [93].

The risk of bowel injuries is higher in laparoscopic surgery for ASBO and bowel resections are significantly more frequent in laparoscopic surgery [72]. In a study the incidence of bowel resection was 53.5 versus 43.4% in laparoscopic versus open procedures [94]. Some authors concluded that laparoscopic approach is safer than the open procedure, but only in the hands of experienced laparoscopic surgeons and in selected patients [95]

Farinella et al reported that predictors for a successful laparoscopic treatment of ASBO are the following:  $\leq 2$  laparotomies in history, appendectomy as the operation in history, no previous median laparotomy incision, unique band adhesion as pathogenic mechanism of small bowel obstruction, early laparoscopic management within 24 h from the onset of symptoms, and no signs of peritonitis on physical examination, and experience of the surgeon [96]

### 2.2.3. Surgical exploration

Surgical exploration tends to search for the site and the cause of obstruction, which can be determined by exploring the transition point between the distended and the flat bowel. Festen [7] reported that in more than 70% of the children with ASBO included in his study, single adhesion was the cause of obstruction, however multi-adhesions, volvulus and compressing abscess were also reported to be found during exploration. The process of exploration is meant to assess the bowel viability which will further guide to choose the right surgical procedure.



**Figure 29:** Perioperative pictures of one of our patients, showing a congenital adhesive band obstructing the small bowel

#### 2.2.4. Assessment of bowel viability [97]

It is a critical process. The evaluation of bowel viability must be done after the removing of the obstruction ( adhesiolysis or untwisting bowel in case of volvulus). Based on several signs this process helps to know if the intestine is going to survive or not :

**Bowel is viable if:**(1) its surface is glistening,(2) its colour is pinkish, or only slightly blue, (3) it feels resilient like normal bowel,(4) it contracts sluggishly (like a worm) when you pinch it, (5) pulsations are seen in the vessels which run over the junction between it and its mesentery

**Bowel is not viable if:**(1) it tends to dry out and its surface is no longer glistening, (2) it is greyish purple, dark purplish-red or even black,(3) it feels like blotting paper, (4) it does not contract when you pinch it,(5) its blood vessels are not pulsating or are clotted.

**If the process of assessment is inconclusive,** remove the cause of thestrangulation, apply a warm, moist pack to it, and wait10mins. If it is viable, its color will change from dusky to normal pink. If this happens, it is alive even if youcannot feel the pulsations of the mesenteric vessels.It may be viable if some areas remain purplish because ofbruising. But if these areas are large, or do not improve incolor, consider all the discolored bowel to be non-viable




**Figure 30:** perioperative picture showing bowel ischemia

2.2.5. Surgical procedures

➤ **Adhesiolysis [97,98]**


surgical adhesiolysis is the process of freeing adhesions, it is the current method of managing adhesions .during this process there is a great danger of bowel perforation. and the safest way to separate adhesions is to use the 'push and spread technique' preferably with Metzenbaum's or McIndoe's scissors, Using the outer sides of the blades to spread the tissues which are not so blunt .



**Figure 31:** illustration images of adhesiolysis [97]

➤ **Retrograde decompression [97,99]**


Is the method of choice, provided the bowel is not too oedematous and friable. It is useful for the entire small bowel. It consist of milking the contents of the entire small bowel between the index and the middle fingers, starting from the jejunum-ileum junction and progressing proximally into the stomach . as the decompression is in process the gastric fluid need to be aspirated through a large bore nasogastric tube working properly to avoid fluid spill which may cause aspiration .



**Figure 32:** image illustrating the process of retrograde decompression[99]

➤ **Enterotomy decompression [97]**

It consists of making a longitudinal opening in the anti-mesenteric border of the bowel, then inserting a decompressor and sucking out gas and fluid. this method is rarely used it can be the best method in case of bowel resection, during which bowel content is drained from the open ends of the bowel before proceeding to resection.



**Figure 33: illustration image of enterotomy decompression[97]**

➤ **Bowel resection and anastomosis [97]**

If a section of bowel is not viable, it must be resected and the two ends of the bowel joined by anastomosis. Though those ends which will be joined must be viable pink bleeding ends, and the bowel must be emptied before anastomosis

usually ends-to-ends anastomosis is performed, joining their serosal surfaces together by 2 layers with an interrupted or continuous suture. At the end of the procedure, the bowel is washed with warm fluid and gently squeezed on either side to test the anastomosis leak. When it is not safe to make an ends-to-ends anastomosis; a stoma is performed.

**Table XVII:the surgical procedures performed by differentauthors**

<b>Surgical procédures</b> <b>Author</b>	<b>Adhesiolysis</b>	<b>Retrograde decompression</b>	<b>Resection – Anastomosis</b>	<b>Enterotomy</b>
<b>N. Erturk [28]</b>	84.5%	not reported	13.3%	2.2
<b>Akgur [18]</b>	82.5%	–	14.6%	2.5
<b>Vijay**[34]</b>	66%	–	33%	27
<b>Eason**[16]</b>	88%	–	9.4%	2.2
<b>our study</b>	<b>93%</b>	<b>93%</b>	<b>7%</b>	<b>0</b>

\*\* Excluding patients managed with immediate surgery

#### 2.2.6. Post-operative outcomes

Many studies have reported the occurrence of postoperative complications [16,17,18].Eason reported a complication rate of 12% [16]. The most common complications observed were surgical site infection, wound dehiscence, prolonged ileus, sepsis and peritonitis after anastomosis leakage, and pneumonia.

We have registred one case of sepsis and peritonitis that occurred after operative management .

## VII. EVOLUTION

### 1. Length of hospital stay

**Table XVIII: the mean length of hospital stay reported by authors**

Author	the mean time of hospital stay ( days )		
	standard conservative management	conservative management with WSCA	operative management
Eson (2010) [16]	6.4	-	immediate = 10.4 Vs Delayed= 14
Feigin [33]	4.2	-	10.2
Nasir [17]	5	-	13
Bonnard [27]	6.5	3	-
Abdelkader [89]	-	3.6	-
F Linden[20]	6.2	3.6	-
our study	2.1	3.5	6.1

The length of hospital stay depends on the therapeutic approach and the complications encountered. The studies reviewed reported a length of hospital stay that ranges from 4.2 to 6.5 days for the patients who were managed conservatively without using WSCA, while this length was shorter for the patients who were managed by WSCA, it ranged between 3 and 3.6 days . in the other hand the operative management required much longer time of hospital stay in comparison to the conservative management , the length of stay for patients treated operatively ranged between 6.1 and 14 days . Nasir [17], Feigin [33] and Eson[16] concluded that conservative management is associated with a shorter length of hospital stay than the operative management. Moreover, the administration of WSCA reduces the length of hospital stay significantly compared to the standard conservative management.

Our study reported a shorter length of hospital stay for the patients managed conservatively than those managed operatively, which is consistent with the literature. Conversely, the length of stay for patients managed with WSCA was a little longer compared to the standard protocol which can be explained by the small number of the patients, the inequality of number between the two groups, and the heterogeneity of the previous surgical conditions.

## **2. Recurrence**

Recurrent ASBO continues to be a concern long after the initial treatment of obstruction, with a reported range of 16-53% [18,31,100,101]. Although surgical treatment of ASBO may lead to the additional formation of adhesions, and recurrent episodes of ASBO. The recurrence is slightly higher for patients treated conservatively than those managed operatively; moreover this rate rises for both approaches after a 5 years period[72]. Akgur[18] reported a double recurrence rate after conservative management compared with the operative management (which was 36.47% vs 18.75%), he also reported that recurrent ASBO episodes could be treated conservatively and the previous treatment has no effect on the rate of success of this management.

No agreement exists about the possibility to predict the recurrence risk[71] .though; some authors reported that Factors associated with a higher risk of recurrence are age < 40 years, matted adhesion and postoperative surgical complications[102], moreover, multiple previous operations were associated with increased risk of ASBO recurrence. Lautz et al[39] reported that patients with 3 or more previous operations had a relative recurrence risk of 3.2

In our study, we encountered 4 cases (6.5%) of ASBO recurrence, 3 were previously managed conservatively, and only one had undergone surgery, which is accordant with studies reports.

## **3. Mortality**

Akgur et al [18] reported a mortality rate of 1.1% (two deaths), one occurred after preoperative perforation complicated to advanced peritonitis and the other after sepsis caused by peritonitis resulting from leakage.

In Festen's study [7] death was encountered in two cases (6.6%), one child died of peritonitis and sepsis caused by an intestinal fistula after decompressive enterotomy, the other one died of sepsis after mesenteric thrombosis.

In our studied group, we have registered two deaths (3.5%). One of them died after 24 hours of conservative management, from a sudden cardiopulmonary arrest secondary to massive inhalation of gastric content. The second one was managed operatively after 72 hours of unsuccessful conservative management; he died from sepsis after being transferred to pediatric critical care medicine.

## VIII. PREVENTION

Enormous effort has been applied to reduce adhesion formation. There are various approaches to prevent those adhesions which can be separated to three main categories: adjusting surgical techniques, Pharmacological Adjuvant Therapy, Adjuvant Barrier Therapy between or over damaged surfaces, those last two approaches are most common and modern [35].

## 1. Surgical techniques :[35 , 1 , 71 ]

Postsurgical coalescing adhesions will form only when both contacting peritoneal surfaces have been traumatized during surgery. There are many effective measures, which minimize peritoneal trauma and lead to less adhesion formation; those measures include :

- pursuing general principles of atraumatic, gentle, and bloodless surgery during either laparoscopy or laparotomy
- careful and delicate handling of the bowel in order to reduce severe trauma, keeping tissues moist with irrigation, and avoiding large abdominal wounds and unnecessary dissection
- using micro and atraumatic instruments to reduce serosal injury
- non closure of the peritoneum during the closure of abdominal wounds, since peritoneal closure may induce ischemia and predisposing the site to decreased fibrinolytic activity and increased adhesion formation.
- Using polyglycolic acid derivatives and monofilament synthetics sutures which are associated with less tissue reaction, whereas catgut suture even rapidly absorbed it is highly reactive
- Areas at higher risk of adhesion development may be covered with omentum, peritoneal flaps, falciform ligament, and broad ligament
- Using powder-free gloves (talc and starch), and avoid contact of the peritoneum with other foreign materials such as gauze lint, sutures, and material extruded from digestive tract because they may be responsible for peritoneal inflammatory reaction which potentiates adhesion formation

- In the presence of intraperitoneal blood, it should be aspirated in irrigation solution and adequate hemostasis must be performed.
- Preferring minimally invasive/laparoscopic surgical techniques, since de novo adhesion formation occurs more frequently in patients undergoing laparotomy.

## **2. Pharmacological adjuvant therapy :[1,35,109]**

A variety of pharmacological agents have been studied including steroidal, nonsteroidal anti-inflammatory drugs, heparin, antihistamines, antibiotics, antioxidants, antiangiogenics, and antifibrotics. Those Drugs may be administered systemically, or ideally, locally with minimal systemic effect, and they are directed against various causes and components of the inflammatory process and/or of adhesion formation.

unfortunately, even some of those drugs have been proven to be effective in animals models. There are obvious implications for biocompatibility and toxicity of any material used for adhesion prevention, furthermore there still no clinical trials that have been done

## **3. Adjuvant Barrier Therapy: [1,35 ,109,110,111]**

Most clinical experience is with intraperitoneal adhesion barriers, applied at the end of surgery with the aim to separate injured peritoneal and serosal surfaces until complete adhesion free healing has occurred. Efficacy of anti-adhesion barriers in open surgery has been well established for reducing the incidence of adhesion formation.

### ➤ **Barrier solution :**


Hyaluronic acid has been combined with Phosphate-Buffered-Saline (HAPBS) into a macromolecular solution to prevent adhesion formation, marked as Separacoat<sup>®</sup>. This solution is applied intraoperatively, prior to dissection, to protect peritoneal surfaces from indirect surgical trauma (e.g., abrasion and desiccation) rather than postoperatively to separate surfaces after they are traumatised. In both animals models and human studies, HAPBS solution was documented to have safely and significantly reduced serosal damage and decreased incidence, extent, and

severity of de novo adhesions.

Adept® is a liquid agent consists of 4% icodextrin in an iso-osmotic electrolyte solution. Icodextrin is high-molecular-weight glucose polymer which results in a prolonged intraperitoneal residence time of the solution allowing abdominal organs to be separated by flotation. The operative handling of Adept is simple as it is introduced via instillation into the peritoneal cavity following laparoscopy where it remains until absorption. It is important to carefully close the abdominal wounds subcutaneously to prevent post-operative leakage of the solution. For this reason, Adept is also not well suited for laparotomy. This solution has demonstrated efficacy in the secondary prevention of adhesions following laparoscopy for adhesiolysis. Unfortunately, Icodextrin has a minimal impact on new adhesion formation.

➤ **Solid synthetic barrier :**

Hyaluronic acid carboxymethylcellulose (HA/CMC) marketed as Seprafilm® is a solid synthetic barrier that prevents denuded epithelial tissues from touching. It has documented efficacy for abdominal and pelvic surgery, and It has been shown to reduce the incidence, extent, severity, and surface area of postsurgical adhesions, in addition to that it was approved by FDA for abdominal and pelvic laparotomy. beside the high price, the limitations of HA/CMC is that it is indicated for open procedures only and can be difficult to handle because the film is brittle and cracks easily. Also, it fixes upon contact with a moist tissue and cannot easily be moved after application which requires very careful handling. However, it is regularly used off-label in laparoscopic surgery using a cigar roll fashion that eases application without reducing its effectiveness.



---

## *CONCLUSION*

---



At the end of this work, and in light of our findings and after analyzing the related studies, we come out with those conclusions :

Adhesive small bowel obstruction is one of the leading emergencies encountered in pediatric surgery that needs early and effective management.

The Peritoneal adhesions are a result of a pathological healing process that follows peritoneal injury, and they occur most commonly after abdominopelvic surgery. However, those adhesions can be congenital.

ASBO can occur at any age, and also after laparotomy for any surgical condition, however history of perforated appendicitis is associated with a high incidence of ASBO

Abdominal pain and vomiting are constant symptoms of ASBO, while obstipation and abdominal distension can be missing in some cases

Abdominal X-ray in orthostatic position plays a crucial role in small bowel obstruction diagnosis. Abdominal CT scan performs better than plain X-ray in identifying the cause of obstruction and recognising complications. Nonetheless, it should not be routinely implemented in the diagnosis making process. The biological tests help to confirm the clinical suspicion of volume depletion and electrolytes imbalances, and also assist recognition of bowel ischemia.


Conservative management, either with standard decompressive therapy or using water-soluble contrast agent, is a safe and effective approach in managing ASBO in selected children without predictor signs of strangulation and bowel ischemia. However, in the absence of clinical indicators suggesting improvement, this approach should not exceed 48 hours of trial.


Laparotomy is used for children with strangulating ASBO and after the failure of conservative management trial. Inappropriate patients, the laparoscopic approach can also be used for managing this condition.

Further prospective studies are needed to establish optimal management for ASBO in the pediatric population.

**Algorithm for the management of ASBO in children :**

**Child admitted for  
ASBO**





---

*APPENDIX*

---



## PATIENT'S MEDICAL SUMMARY

**I- IDENTITY :**

- Entry code : .....
- Full Name : .....
- Age : .....
- Gender : .....
- Date of admission : .....
- Date of discharge : .....

**II- PAST MEDICAL HISTORY**

1) **MEDICAL HISTORY:** NO  YES  (precise )

.....

2) **SURGICAL HISTORY:** NO  YES  (precise )

A) The Condition

- Appendicitis with perforation  Appendicitis without perforation   
 Intussusception  Malrotation  Hirschprung's disease   
 Other  (precise) .....

B) type of intervention ( Laparotomy  Laparoscopy  )

C) site of intervention ( supramesocolic space  inframesocolic space  )

**III- HISTORY OF PRESENTING COMPLAINT (HPC)**

1) **Symptoms**

A- **Abdominal pain :** NO  Yes  (if so, precise the abdominal region)  
 ( Right iliac  Right lumbar  left lumbar  Epigastric  Umbilical   
 hypogastrium  Right hypochondrium  Left hypochondrium  Diffuse  )

B- **Vomiting :** NO  YES   
 If so , precise the character ( non-bilious  Bilious  )

C- **OBSTIPATION (Failure to pass flatus and stool):** NO  YES

D- **Other symptoms :** .....

2) **TIME TO ONSET OF SYMPTOMS AFTER THE FIRST SURGERY**

- A- 0 – 6 Weeks  B- 7weeks – 3months  C- 3months - 12 months   
 D- 13 months – 5 years  E – more than 5 years



**VI- TREATMENT**

**1- CONSERVATIVE MANAGEMENT**

- A- Usual Conservative protocol** NO  YES
- Nasogastric aspiration  Analgesics  Antispasmodics
- Antiemetics  Antibiotics
- Resuscitation with intravenous fluids and electrolytes,

a) **Duration of the usual conservative protocol :**.....

- b) Treatment outcome :** Successful  Unsuccessful
- (.....)

- B- Water-soluble contrast ( Gastrografin ) protocole** NO  YES

a) **The outcome :** Successful  Unsuccessful

**2- OPERATIVE MANAGEMENT** NO  YES

If so , precise :

- A- Timing :** EARLY (INITIALLY)  DELAYED(after conservative management)
- If DELAYED ,pricise the time elapsed after the admission.....

**B- Surgical Approach**

Laparotomy  laparoscopy

**C- Surgical exploration :**.....

**D- Operative procedure**

Adhesiolysis  Retrograde decompression  resection of bowel

decompressive enterotomy  perforation repair

Other  (.....)

**E- Postoperative outcomes**

- Normal with No complication
- Complication  ( wound abcess  fistula  ventral hernia
- Dehiscence  Pneumonia  other  (.....)

**VII- Evolution**

- 1) **Lenght of hospital staying** .....days
- 2) **Death** NO  YES
- 3) **Recurrence** NO  YES

## Appendix II:

### Anatomical and physiological Reminder

#### 1. Anatomical Reminder [104,105]


##### 1.1. The small bowel

The small intestine is the longest part of the gastrointestinal tract with approximately 6 to 7 meter long .it extends from the pyloric orifice of the stomach to the ileocecal fold, and it consists of the duodenum, the jejunum, and the ileum .

##### 1.1.1. Duodenum

Duodenum is first part of the small intestine, is about 20 - 25 cm long and its lumen is the widest of the small intestine . this C-shaped structure is retroperitoneal except for its beginning , and is adjacent to the head of the pancreas . it is divided into four parts and terminates at the duodenojejunal flexure.

The arterial supply to the duodenum includes superior pancreaticoduodenal branches of the gastroduodenal artery and the inferior pancreaticoduodenal branch of the superior mesenteric artery , those two arteries anastomose with each other .the veins draining the duodenum follow the arterial supply and terminate in the portal venous system.



**Figure 34 : anterior view of the duodenum with anterior wall opened**

### 1.1.2. Jejunum


By convention, the proximal two-fifths represent the jejunum and the remainder the ileum. However, no precise anatomical feature marks the junction between the two, there being a gradual morphological transition along the whole length of the small intestine. The jejunum is larger in diameter and has a thicker wall than the ileum. In addition to that the inner mucosal lining of the jejunum is characterized by numerous prominent folds that circle the lumen (plicae circulares) but diminish in both size and number along the intestine so that in the distal ileum they are often absent.

The arterial supply to the jejunum includes jejunal arteries from the superior mesenteric artery.


### 1.1.3. Ileum

The ileum makes up the distal three-fifths of the small intestine. Compared to the jejunum, the ileum has thinner walls, fewer and less prominent mucosal folds (plicae circulares) and it contains lymphoid tissue arranged in discrete clumps (Peyer's patches). The ileum opens into the large intestine where the cecum and ascending colon join together and it projects two flaps into the lumen of the large intestine (the **ileocecal fold**). Additionally, musculature from the ileum continues into each flap, forming a sphincter which prevents reflux from the cecum to the ileum, and regulates the passage of contents from the ileum to the cecum.

The arterial supply of the ileum comes from ileal arteries of the superior mesenteric artery and an ileal branch of the ileocolic artery (from the superior mesenteric artery).



**Figure 35: Wall structure of the jejunum and ileum**




**Figure 36: Anterior view showing the arterial supply of small bowel**

## 1.2. The large intestine

The large intestine extends from the distal end of the ileum to the anus. It is approximately 1.5 m long in adult, and its lumen is more wide than the small intestine. It comprises the caecum, appendix, colon, rectum and anal canal. The caecum and appendix lie in the right iliac fossa, while the colon runs a circuitous course forming ascending colon, right colic flexure, transverse colon, left colic flexure, the descending colon and the sigmoid colon, before continuing on posterior wall the pelvic cavity as the rectum.

The large intestine contains Sacculations called (the haustra of the colon) and segregation of longitudinal muscle in its walls into three narrow bands (the taeniae coli).

The blood supply of the caecum, appendix, ascending colon and most of the transverse colon is provided by the superior mesenteric vessels. The remainder of the colon is supplied by the inferior mesenteric vessels. The venous drainage follows the arterial supply and terminates in the portal venous system.




**Figure 37:** the arterial supply of the large bowel


### 1.3. The peritoneum and peritoneal cavity

The peritoneum is a serous membrane with parietal and visceral layers. It has almost the same surface area as the epidermis which is approximately between 1 to 2 m<sup>2</sup>. The parietal peritoneum lines the walls of the cavity and the visceral peritoneum covers the viscera. Between the two layers of peritoneum there is a potential space (the peritoneal cavity). Usually, this cavity contains only a few millilitres (mL) of serous fluid. It comprises the greater and lesser sacs. The greater sac is very extensive; it lies between the diaphragm above into the pelvic cavity below. While the lesser sac or the omental bursa is located in the upper part of the abdomen, behind the stomach and communicates with the greater sac through a narrow opening (the omental or epiploic foramen). The greater sac is divided by the transverse mesocolon into supracolic and infracolic compartments

The structure of the peritoneum consists of a single outer layer of mesothelial cells that are supported by a basement membrane and that detach readily with even the slightest trauma. The submesothelial layer consists of components of the extracellular matrix, along with capillaries and lymphatics. Fluid resorption and diffusion occurs freely across these layers. The fluid in the peritoneal cavity contains several different cell types, including leukocytes and macrophages. These cells, along with the mesothelium, secrete various cellular mediators that have roles in peritoneal healing, enabling modulation of the inflammatory response over a large surface area.



**Figure 38:** Illustration image showing the disposition of peritoneum layers in the peritoneal cavity



**Figure 39:** Midsagittal section of the abdominopelvic cavity showing the disposition of peritoneum

## 2. Physiological reminder [104,106,107]

### 2.1. Physiology of the digestive system

The primary function of the digestive system is to transfer nutrients, water, and electrolytes from the food we eat into the body's internal environment. There are four basic digestive processes: motility, secretion, digestion, and absorption :

- **Motility:** refers to the muscular contractions that mix and move forward the contents of the digestive tract. It helps to mix food with the digestive juices, which promotes the digestion of the food and facilitates absorption by exposing all parts of the intestinal contents to the absorbing surfaces of the digestive tract.
- **Secretion:** Several digestive juices are secreted into the digestive tract lumen by exocrine glands along the route, each with its own specific secretory product. They consist of water, electrolytes, and specific organic constituents important in the digestive process. Furthermore there are endocrine cells located in the digestive tract wall secrete gastrointestinal hormones into the blood that help control digestive motility and exocrine gland secretion
- **Digestion:** refers to the biochemical breakdown of the structurally complex foodstuffs of the diet into smaller, absorbable units by the enzymes produced within the digestive system :
  - ✓ Carbohydrates are degraded into simple sugars or monosaccharides ("one-sugar" molecules), such as glucose, fructose, and galactose. These monosaccharides are the absorbable units for carbohydrates.
  - ✓ Proteins are degraded primarily into their constituent amino acids as well as a few small polypeptides, both of which are the absorbable units for protein.
  - ✓ Triglycerides, which are neutral fats consisting of a glycerol with three fatty acid molecules attached. During digestion two of the fatty acid molecules are split off, leaving a monoglyceride and free fatty acids which are the absorbable units of fat.

- **Absorption** : Through the process of absorption, the small absorbable units that result from digestion are transferred from the digestive tract lumen into the blood or lymph .water and all products of carbohydrate, protein, and fat digestion, as well as most of the ingested electrolytes, and vitamins are mostly absorbed in duodenum and jejunum, very little occurs in the ileum . Otherwise, vitamin B12 and bile salts are exceptionally absorbed in Ileum, because the specialized transport mechanisms for these two substances are located only in this region.

The colon extracts more H<sub>2</sub>O and salt from the contents. What remains to be eliminated is known as feces.

## 2.2. The functions of the peritoneum

- **Movement of viscera**: the serous fluid secreted by the peritoneum allows free movement of abdominal viscera and peristalsis of the stomach and intestine
- **Defense and Protection**: it guards against germs by phagocyte and plays a role in the immunological response by activating lymphocytes and monocytes .also the greater omentum has a particular power to move toward the site on infection .it plays the role of “abdominal policeman”
- **Absorption**: it absorbs fluid effusion to blood capillaries
- **Repair**: the mesothelial cells of the serosa synthesize inflammatory cytokines, growth factors, and extracellular matrix (ECM), which are actively involved in serosal wound healing and the regulation of peritoneal inflammation.



---

# *ABSTRACT*

---



### **Abstract:**

Adhesive small bowel obstruction (ASBO) is one of the leading causes of surgical emergencies. Unfortunately, it does not spare children.

This is a retrospective Study involved 61 cases of ASBO in 57 children, admitted in pediatric surgery department of the Mohamed VI Teaching Hospital of Marrakech, within a period of 7 years, between January 2012 and December 2018. It aims to review the epidemiological, clinical, and paraclinical features, and also outlines the therapeutic aspects of this condition.

A masculine predominance of 80.7% was found. The mean age of our patients was 7 years and 5 months with extremes ranging between 8 days and 15 years. Perforated appendicitis was the most common previous surgical condition leading to ASBO. Three patients (5.26%) have no medical nor surgical history, and one patient was treated for abdominal tuberculosis. Pain and vomiting were present in all children. Obstipation was found in 97% of cases. abdominal radiographs showed air-fluid levels and distended loops of bowel in all the patients.

11 cases were operated-on immediately, and 50 cases have had non-operative management trial, using either standard nasogastric decompression or Water-soluble contrast agent, which was successful in thirty-four cases (68%), the remaining 14 cases with unsuccessful conservative management underwent delayed surgery. The surgical management consisted of adhesiolysis with retrograde decompression in 23 patients and bowel resection in 2 patients. The mean length of hospital stay was 3 days for the patients managed conservatively and 6.1 days for the patients who underwent surgery. Recurrence occurred in 4 patients. Mortality was encountered in two patients (8%). Conservative management is a safe and effective therapeutic approach for managing ASBO in selected children without predictor signs of strangulation and bowel ischemia.

## Résumé :

L'occlusion intestinale par bride (OIB) est l'une des principales urgences chirurgicales, malheureusement, elle n'épargne pas les enfants.

ceci est une étude rétrospective qui a inclus 61 cas de OIB chez des enfants qui ont été admis au service de chirurgie pédiatrique de l'hôpital Universitaire MED VI à Marrakech, sur une période de 7 ans, entre Janvier 2012 et Décembre 2018. cette étude a pour objectif d'étudier les aspects épidémiologique, clinique et paraclinique et aussi de souligner les aspects thérapeutiques de cette pathologie.

Une prédominance masculine de 80.7% a été retrouvée chez nos patients. l'âge moyen a été 7ans et 5 mois avec des extrêmes entre 8 jours et 15 ans. l'appendicite perforée était l'antécédent chirurgical le plus fréquent. trois patients (5.26%) n'avaient aucun antécédents médicaux ou chirurgicaux et un patient a été suivi pour tuberculose abdominale. la douleur abdominale et les vomissements étaient présents chez tous nos patients, et l'arrêt des matières et des gaz dans 97% des cas. les radiographies abdominale ont montré des niveau hydro-aérique et la dilatation des anses intestinales chez tous les patients.

11 cas ont été opérés immédiatement, et 50 cas ont bénéficiés de l'essai du traitement conservateur soit par l'aspiration nasogastrique standard ou l'administration de produit de contraste hydrosoluble, cette essai a été efficace chez 34 cas (68%), les 14 cas restants dont le traitement conservateur a échoué ont subi d'une exploration chirurgicale retardée. le traitement chirurgical a consisté l'adhésiolyse avec l'entérovidange rétrograde chez 23 patients et la résection intestinale chez 2 patients. la moyenne de séjour à l'hôpital était 3 jours pour les patients traités conservativement et 6.1 jours pour les patients qui ont subi la chirurgie. la récurrence a été enregistrée chez 4 patients. la mortalité est survenue chez deux patients (8%). le traitement conservateur est un moyen thérapeutique sûr et efficace pour la prise en charge de OIB chez des enfants bien sélectionnés sans signes prédictifs de strangulation et d'ischémie intestinale.

## ملخص

يعتبر انسداد المعيد الدقيق بالاتصاقات أحداً هاماً للمستعجلات الجراحية ، للأسف هذه الحالة المرضية لاتعفيا لاطفال .

هذه دراسة استرجاعية شملت 61 حالة لانسداد المعيد الدقيق بالاتصاقات تلدى 57

طفلاً استقبلوا في مصلحة جراحة الاطفال بالمستشفى الجامعي محمد السادس بمرآكش ، خلال مدة 7 سنوات بينيناير 2012 و دجنبر 2018 . وترمي هذه الدراسة المراجعة المميز اتالوبائية ، والسريرية ، والاسريرية ، وكذا الكبار از مختلفا لوجهالعلاجية لهذه الحالة المرضية .

تم ملاحظة سيادة العنصر الذكوري ( بنسبة 80.7 % ) . متوسط العمر لدمرضانا كان 7 سنوات و 5 اشهر ومختلفا لعمار مناصغر ها لأكبر ها كان بين 8 ايام و 15 سنة . ثلاث مرضى ( 5.26% ) لم تكن لديهما أية سوابق طبية او جراحية ، مريض واحد كان يتابع علاجاً لمرض سلال البطن .

مرضات لها بالزائدة الدودية كانتا الحالة الجراحية السابقة المؤدية اكثر الانسداد المعيد الدقيق بالاتصاقات . الامواتقينا كان متواجداً عند كلا لاطفال ، فيحين لا مساك المعند و جدلى 97% من الحالات . الاشعة السينية للبطن بينت وجود مستويات تبين الهواء - السائل وكذا الوجود عروا تامعاء منتفخة عند كلا المرضى .

11 حالة استندت عند خلع جراحيفوري ، و 50 حالة استفادوا من المعالجة التحفظية التجريبية ، باستعمال تخفيف الضغط الانفي بالمعدي المعياريا واستعمال تباينيدو ابيالماء . المعالجة التحفظية كانت ناجحة عنداربوعو ثلاثون حالة (68% ) ، الاربعة عشر حالة المتبقية التليمتفاحمعها المعالجة التحفظية خضعو للجراحة المؤجلة .

المعالجة الجراحية تتضمنت عملية فكالاتصاقات وتفرغ الامعاء الرجوعيلدى 23 مريضو عملية قطع الامعاء لدمرضين . متوسط مدة الاقامة في المستشفى هو 3 ايام بالنسبة للمرضى المعالجينو فالمعالجة التحفظية ، و 6.1

يوم بالنسبة للمرضى الذين خضعو للجراحة . ايبوبة المرضى حدثت لدى 4 مرضى . الوفاة سجلت لدمرضين ( 8% ) . المعالجة التحفظية هي طريقة امنة وفعالة لمعالجة انسداد المعيد الدقيق بالاتصاقات عند بعض الاطفال الامنتقيدون علاما تنبؤوا باختناق ونقصار تواء الامعاء .



---

# *BIBLIOGRAPHY*

---



1. **Theodoros Liakakos, Nikolaos Thomakos & al.**  
Peritoneal Adhesions: Etiology, Pathophysiology, and Clinical Significance.  
Dig Surg 2001;18:260–273.
2. **M. H. Alwan, A. M. van Rij, and S. F. Greig.**  
Postoperative adhesive small bowel obstruction: the resources impacts  
The New Zealand Medical Journal, 1999 Nov 12;112(1099):421–3..
3. **Attard JA, MacLean AR.**  
Adhesive small bowel obstruction: epidemiology, biology and prevention.  
Can J Surg 2007;50:291–300.
4. **Aguayo P, Ho B, Fraser JD, et al.**  
Bowel obstruction after treatment of intra–abdominal tumors.  
Eur J Pediatr Surg 2010;20:234–6.
5. **El–Gohary Y, Alagtal M, Gillick J.**  
Long–term complications following operative intervention for intestinal malrotation: a 10–  
year review.  
Pediatr Surg Int 2010;26:203–6 .
6. **Grant HW, Parker MC, Wilson MS, et al.**  
Population–based analysis of the risk of adhesion–related readmissions after abdominal  
surgery in children.  
J Pediatr Surg 2006;41:1453–6.
7. **Festen C.**  
Postoperative small bowel obstruction in infants and children.  
Ann Surg 1982;196:580–3
8. **Jaime Shalkow, MD , Carmen Cuffari, MD .**  
Pediatric Small–Bowel Obstruction.  
emedicine.medscape.com , Updated: Dec 11, 2015.

9. **Lin L-H, Lee C-Y, Hung M-H, et al.**  
Conservative treatment of adhesive small bowel obstruction in children: a systematic review.  
BMJ Open 2014;4:e005789.doi:10.1136/bmjopen-2014-005789
  
10. **Williams SB, Greenspon J, Young HA, et al.**  
Small bowel obstruction: conservative vs. surgical management.  
Dis Colon Rectum 2005;48:1140-6.
  
11. **Seror D, Feigin E, Szold A, et al.**  
How conservatively can postoperative small bowel obstruction be treated?  
Am J Surg 1993;165:121-5; discussion 125-6.
  
12. **Fevang BT, Jensen D, Svanes K, et al.**  
Early operation or conservative management of patients with small bowel obstruction?  
Eur J Surg 2002;168:475-81.
  
13. **Joseph S. Janik, Sigmund H. Ein et al .**  
An assessment of the surgical treatment of adhesive small bowell obstruction in infants and children.  
Journal of pediatric surgery. 1981; 16: 225-229
  
14. **Timothy B Lautz, MD, Mehul V Raval, M et al .**  
Adhesive Small Bowel Obstruction in Children and Adolescents: Operative Utilization and Factors Associated with Bowel Loss.  
Journal of the American college of surgeons. May 2011, Vol.212(5): 855-861
  
15. **Ahmed H . A I-Salem and Mohammad Oquaish.**  
Adhesive Intestinal Obstruction in Infants and Children: The Place of Conservative Treatment.  
Int Scholarly Research Network. ISRN Surgery Volume 2011, Article ID 645104, 4 pages
  
16. **Gareth A. Eeson, Paul Wales, James J. Murphy**  
Adhesive small bowel obstruction in children: should we still operate ?  
Journal of Pediatric Surgery (2010) 45, 969- 974.

**17. Abdulrasheed A. Nasir, Lukman O. Abdur-Rahman et al.**

Is non-operative management still justified in the treatment of adhesive small bowel obstruction in children?.

African Journal of Paediatric Surgery. JulySeptember2013/Vol 10/Issue 3.

**18. Akgür FM, Tanyel FC, Büyükpamukçu N, et al.**

Adhesive small bowel obstruction in children: the place and predictors of success for conservative treatment.

J Pediatr Surg 1991;26:37-41

**19. Lee CY, Hung MH, Lin LH, et al.**

Evaluation of a water-soluble contrast agent for the conservative management of adhesive small bowel obstruction in pediatric patients.

J Pediatr Surg 2014;50:581-5

**20. Linden AF, et al.**

Evaluation of a water-soluble contrast protocol for nonoperative management of pediatric adhesive small bowel obstruction.

J Pediatr Surg (2018), <https://doi.org/10.1016/j.jpedsurg.2018.10.002>

**21. BOUHNOUN ZAHRA**

L'OCCLUSION INTESTINALE SUR BRIDES POST-OPÉRATOIRE CHEZ L'ENFANT (A PROPOS DE 43 CAS)

These d'obtention du DOCTORAT EN MEDECINE , N° 246/16 , fes 2016

**22. G K. Kouadio , HT TURQUIN.**

Prise en charge des occlusions post-opératoires du grêle par brides et adhérences au CHU de Treich ville à Abidjan.

Médecine d'Afrique Noire ,2004 - 51 (12).

**23. Hiki N, Takeshita Y, Kubota K, et al .**

A seasonal variation in the onset of postoperative adhesive small bowel obstruction is related to changes in the climate.

Dig liver Dis 2004; 36 .

24. **J. Kossi, P. Salminen , et al.**  
The epidemiology and treatment patterns of postoperative adhesion induced intestinal obstruction in varsinais-suomi Hospital District.  
Scandinavian journal of surgery 93: 68 – 00, 2004
25. **Zerey M, Sechrist CW, Kercher KW, Sing RF, Matthews BD, Heniford BT.**  
Laparoscopic management of adhesive small bowel obstruction.  
Am surg 2007; 73 (8): 3 – 8 .
26. **Duron JJ, Silva NJ, Du Montcel ST, Berger A, Muscari F, Hennet H, Veijrieres M, Hay JM.**  
Adhesive postoperative small bowel obstruction : Incidence and risk factors of recurrence after surgical treatment: a multicenter prospective study. Ann surg. 2006; 244 (5): 750 – 7.
27. **Bonnard A, Kohaut J, Sieurin A, et al.**  
Gastrografen for uncomplicated adhesive small bowel obstruction in children.  
Pediatr Surg Int 2011;27:1277-81.
28. **Erturk N, Sonmez K, Kale N.**  
Evaluation of a series of patients operated for adhesive intestinal obstructions.  
Int J Surg Med. 2017; 3(3): 167-172. doi:10.5455/ijsm.adhesions-abdominal-surgeryinfants
29. **Chirdan LB, Uba AF, Igoche MI, Soo CT, Osagie OO**  
small intestinal obstruction from peritoneal adhesions in children.  
original scientific articles.
30. **ROHR S.**  
Occlusions intestinales aiguës.  
Faculté de Médecine ULP Strasbourg Année 2002. Item 217. Module 11.
31. **Young, J.Y., et al.**  
High incidence of postoperative bowel obstruction in newborns and infants.  
J Pediatr Surg, 2007. 42(6): p. 962-5; discussion 965.

32. **X.Y. Cao<sup>1</sup>, J. Jiang<sup>2</sup> \*, M.-S. Jin<sup>3</sup>, L. He<sup>1</sup>, Q. Wang<sup>1</sup>, J. Suo .**  
Treatment for abdominal tuberculosis caused adhesive small bowel obstruction using ileus tube combined somatostatin.  
Poster Presentations, Poster Session Tuberculosis , presented at the 5th Ditan International Conference on Infectious Diseases: Infectious Diseases in the Resistance Era , Beijing, China
33. **Elad Feigin et al .**  
The 16 golden hours for conservative treatment in children with postoperative small bowel obstruction.  
Journal of Pediatric Surgery (2010) 45, 966–968 .
34. **Vijay, MCh, C Anindya, MCh, P Bhanu et al .**  
Adhesive Small Bowel Obstruction in Children – Role of Conservative Management.  
Med J Malaysia Vol 60 No 1 March 2005 .
35. **Allison H. Maciver, Michael McCall, A.M. James Shapiro.**  
Intra–abdominal adhesions: Cellular mechanisms and strategies for prevention  
International Journal of Surgery 9 (2011) 589e594
36. **Weibel MA, Majno G.**  
Peritoneal adhesions and their relation to abdominal surgery.A postmortem study.  
Am J Surg. 1973;126(3):345–53.
37. **Butt MU, Velmahos GC, Zacharias N, Alam HB, de Moya M, King DR.**  
Adhesional small bowel obstruction in the absence of previous operations: management and outcomes.  
World J Surg. 2009;33(11):2368–71.
38. **KH. Yang et al**  
Congenital adhesion band causing small bowel obstruction: What’s the difference in various age groups, pediatric and adult patients?  
BMC Surgery (2016) 16:79 ,DOI 10.1186/s12893–016–0196–4.

- 39. Timothy B. Lautz MD, Katherine A. Barsness MD, MS.**  
Adhesive Small Bowel Obstruction – acute management and treatment in children.  
Seminars in Pediatric Surgery, <http://dx.doi.org/10.1053/j.sempedsurg.2014.06.006>
- 40. Andersson REB.**  
Small bowel obstruction after appendicectomy.  
Br J Surg 2001;88:1387– 92.
- 41. Van Eijcka FC, Wijnenb RM, van Goora H.**  
The incidence and morbidity of adhesions after treatment of neonates with gastroschisis and omphalocele: A 30–year review.  
J Pediatr Surg 2008;43:479–83.
- 42. Choudhry MS, Grant HW.**  
Small bowel obstruction due to adhesions following neonatal laparotomy.  
Pediatr Surg Int 2006;22:729–32.
- 43. Jo–Anne P. Attard, MD; Anthony R. MacLean, MD**  
Adhesive small bowel obstruction: epidemiology, biology and prevention .  
Can J Surg, Vol. 50, No. 4, August 2007
- 44. Cynthia A. Gingalewski , Jay Grosfeld , James A.O'Neill Jr ,Eric W. Fonkalsrud , Arnold G.Coran**  
.  
Other Causes of Intestinal Obstruction .  
Chapter 85 . Book of pediatric surgery , Sixth Edition Volume 1
- 45. Borie F., Guillon F., Aufort S.**  
Occlusions intestinales aiguës de l'adulte : diagnostic.  
EMC (Elsevier Masson SAS, Paris), Gastro–entérologie, 9–044–A–10, 2009.
- 46. Lily E. Johnston , John B. Hanks .**  
Small Bowel Obstruction  
Chapter 72 , p842–p850,

- 47. de Vaumas C., Montravers P., Dupont H.**  
Syndromes occlusifs.  
EMC (Elsevier Masson SAS, Paris), Anesthésie-Réanimation, 36-726-A-50, 2008
- 48. R.T. Beyene et al.**  
Intra-abdominal adhesions: Anatomy, physiology, pathophysiology, and treatment . Current Problems in Surgery 52 (2015) 271-319.
- 49. Kasper, Dennis L. et al.**  
Acute intestinal obstruction , Harrison's Principles of Internal Medicine. 19th edition. McGraw Hill Education, 2015 , Chapter 355, Page 1984 .
- 50. Lappas JC, Reyes BL, Maglinte DD.**  
Abdominal radiography findings in small-bowel obstruction: relevance to triage for additional diagnostic imaging.  
AJR Am J Roentgenol 2001;176:167-74
- 51. Thompson WM, Kilani RK, Smith BB, et al.**  
Accuracy of abdominal radiography in acute small-bowel obstruction: does reviewer experience matter?  
Am J Roentgenol. 2007;188(3):W233-W238. doi:10.2214/ AJR.06.0817.
- 52. Craig Hacking , Jeremy Jones et al.**  
Small bowel obstruction .  
rID: 6158 , <https://radiopaedia.org/articles/small-bowel-obstruction?lang=us>
- 53. Suri S, Gupta S, Sudhakar PJ.**  
Comparative evaluation of plain films, ultrasound and CT in the diagnosis of intestinal obstruction  
Acta Radiol 1999;40:422-8

- 54. Chao A, Gharahbaghian L.**  
Tips and Tricks: Clinical Ultrasound for Small Bowel Obstruction – A Better Diagnostic Tool?  
ACEM newsletter. Available from: <http://www.acep.org/Content.aspx?ID=100218>
- 55. Hefny AF, Corr P, Abu-zidan FM.**  
The role of ultrasound in the management of intestinal obstruction.  
J Emerg Trauma Shock. 2012;5(1):84-6.
- 56. Jacob Avila, MD, RDMS**  
Small bowel obstruction, Ultrasound of the Week  
<https://www.ultrasoundoftheweek.com/uotw-20-answer/>
- 57. Mourad Boudiaf, MD.**  
CT Evaluation of Small Bowel Obstruction1 .  
EDUCATION EXHIBIT. 614 May-June 2001. RG f Volume 21 , Number 3
- 58. Mullan CP, Siewert B, Eisenberg RL.**  
Small bowel obstruction.  
AJR Am J Roentgenol 2012; 198: W105-W117 .
- 59. Jaffe TA, Martin LC, Thomas J, Adamson AR, DeLong DM, Paulson EK.**  
Small-bowel obstruction: coronal reformations from isotropic voxels at 16-section multi-detector row CT.  
Radiology 2006; 238: 135-142 [PMID: 16293807 DOI: 10.1148/radiol.2381050489].
- 60. Catel L, Lefèvre F, Laurent V. Canard L, Bresier L, Guillemin FD et al.**  
Occlusion sur bride. Quels critères scanographiques de gravité rechercher ?  
J radiol 2003 ; 84:27-31
- 61. Case courtesy of Dr Ian Bickle.**  
Radiopaedia.org, rID: 48397  
<https://radiopaedia.org/cases/adhesional-small-bowel-obstruction-1>

**62. Case courtesy of Dr Ian Bickle,**

Radiopaedia.org, rID: 29769

<https://radiopaedia.org/articles/small-bowel-faeces-sign-2?lang=us>.

**63. Case courtesy of Dr Michael P Hartung.**

Radiopaedia.org, rID: 59269

<https://radiopaedia.org/cases/closed-loop-small-bowel-obstruction-adhesions-1?lang=us>.

**64. P. Taourel, C. Alili, E. Pages, F. Curros Doyon, I. Millet**

Mechanical occlusions: diagnostic traps and key points of the report

Diagn Interv Imaging, 94 (2013), pp. 805–818.

**65. Case courtesy of Dr Vikas Shah .**

Radiopaedia.org, rID: 55559 .

<https://radiopaedia.org/articles/intramural-bowel-gas?lang=us>.

**66. Trésallet C, Lebreton N, Royer B, Leyre P, Godiris–Petit G, Menegaux F.**

Improving the management of acute adhesive small bowel obstruction with CT–scan and water–soluble contrast medium: a prospective study.

Dis Colon Rectum 2009; 52: 1869–1876 [PMID:

**67. Zielinski, M. D., & Bannon, M. P. (2011).**

Current Management of Small Bowel Obstruction.

Advances in Surgery, 45(1), 1–29.doi:10.1016/j.yasu.2011.03.017

**68. Eren T, Boluk S, Bayraktar B, et al.**

Surgical indicators for the operative treatment of acute mechanical intestinal obstruction due to adhesions.

*Ann Surg Treat Res.* 2015;88(6):325–333. doi:10.4174/astr.2015.88.6.325.

**69. Evennett NJ, Petrov MS, Mittal A, Windsor JA.**

Systematic review and pooled estimates for the diagnostic accuracy of serological markers for intestinal ischemia.

World J Surg. 2009;33:1374–1383.

- 70. van Oudheusden TR, Aerts BA, de Hingh IH, Luyer MD.**  
Challenges in diagnosing adhesive small bowel obstruction.  
World J Gastroenterol. 2013;19(43):7489–7493. doi:10.3748/wjg.v19.i43.7489.
- 71. Catena F, Di Saverio S, Coccolini F, et al.**  
Adhesive small bowel adhesions obstruction: Evolutions in diagnosis, management and prevention.  
World J Gastrointest Surg. 2016;8(3):222–231. doi:10.4240/wjgs.v8.i3.222.
- 72. ten Broek et al.**  
Bologna guidelines for diagnosis and management of adhesive small bowel obstruction (ASBO): 2017 update of the evidence–based guidelines from the world society of emergency surgery ASBO working group.  
World Journal of Emergency Surgery (2018) 13:24 ,<https://doi.org/10.1186/s13017-018-0185-2>.
- 73. Osarumwense David Osifo and Mike Efe Ovueni.**  
Is nonoperative management of adhesive intestinal obstruction applicable to children in a resource–poor country?  
African Journal of Paediatric Surgery , May–August 2010 / Vol 7 / Issue 2.
- 74. Stewardson RH, Bombeck CT, Nyhus LM.**  
Critical operative management of small bowel obstruction.  
Ann Surg 1978;187:189–93.
- 75. Di Saverio S, Coccolini F, Galati M, Smerieri N, Biffi WL, Ansaloni L, et al. Bologna.**  
guidelines for diagnosis and management of adhesive small bowel obstruction (ASBO): 2013 update of the evidence–based guidelines from the world society of emergency surgery ASBO working group.  
World J Emerg Surg. 2013;8(1):42
- 76. Loftus T, Moore F, VanZant E, Bala T, Brakenridge S, Croft C, et al.**  
A protocol for the management of adhesive small bowel obstruction.  
J Trauma Acute Care Surg. 2015;78(1):13–9.

**77. Schraufnagel D, Rajae S, Millham FH.**

How many sunsets? Timing of surgery in adhesive small bowel obstruction: a study of the nationwide inpatient sample.

J Trauma Acute Care Surg. 2013;74(1):181-7

**78. Keenan JE, Turley RS, McCoy CC, Migaly J, Shapiro ML, Scarborough JE.**

Trials of nonoperative management exceeding 3 days are associated with increased morbidity in patients undergoing surgery for uncomplicated adhesive small bowel obstruction.

J Trauma Acute Care Surg. 2014;76(6): 1367-72.

**79. Hok-Kwok Choi, FRCS (Edin), et al .**

Therapeutic Value of Gastrografin in Adhesive Small Bowel Obstruction After Unsuccessful Conservative Treatment A Prospective Randomized Trial .

ANNALS OF SURGERY .Vol. 236, No. 1, 1-6.

**80. Skucas J.**

Anaphylactoid reactions with gastrointestinal contrast media.

AJR Am J Roentgenol 1997; 168:962-964.

**81. Trulzsch DV, Penmetsa A, Karim A, et al.**

Gastrografin-induced aspiration pneumonia: a lethal complication of computed tomography.

South Med J 1992; 85:1255-1256..

**82. Fevang BT, Jensen D, Fevang J, et al.**

Upper gastrointestinal contrast study in the management of small bowel obstruction: a prospective randomised study.

Eur J Surg 2000;166:39-43.

**83. Abbas S, Bissett IP, Parry BR.**

Oral water soluble contrast for the management of adhesive small bowel obstruction.

Cochrane Database Syst Rev 2007;18(3): CD004651.

**84. Haule C, Ongom PA, Kimuli T.**

Efficacy of gastrografen compared with standard conservative treatment in management of adhesive small bowel obstruction at Mulago National Referral Hospital.

J Clin Trials 2013;3(4) [pii: 1000144].

**85. Assalia, A., et al.**

Therapeutic effect of oral Gastrografen in adhesive, partial small-bowel obstruction: a prospective randomized trial.

Surgery, 1994. 115(4): p. 433-7.

**86. Di Saverio, S., et al.**

Water-soluble contrast medium (gastrografen) value in adhesive small intestine obstruction (ASIO): a prospective, randomized, controlled, clinical trial.

World J Surg, 2008. 32(10): p. 2293-304.

**87. Branco BC, Barmparas G, Schnüriger B, Inaba K, Chan LS, Demetriades D.**

Systematic review and meta-analysis of the diagnostic and therapeutic role of water-soluble contrast agent in adhesive small bowel obstruction.

Br J Surg 2010;97:470-8.

**88. Atahan K, Aladagli I, Cökmez A, et al.**

Hyperosmolar water-soluble contrast medium in the management of adhesive small-intestine obstruction.

J Int Med Res 2010; 38(6):2126-34.

**89. Hesham Abdelkader, et al .**

Gastrografen in the management of adhesive small bowel obstruction in children: a pilot study.

Annals of Pediatric Surgery 2011, 7:3-6.

**90. Nordin A, Freedman J.**

Laparoscopic versus open surgical management of small bowel obstruction: an analysis of clinical outcomes. Surg Endosc. 2016;30(10):4454-63

**91. Szomstein S, Lo ME, Simpfendorfer C, Zundel N, Rosenthal RJ.**

- Laparoscopic lysis of adhesions.  
World J Surg. 2006;30(4):535-40
- 92. Hackenberg T, Mentula P, Leppaniemi A, Sallinen V.**  
Laparoscopic versus open surgery for acute adhesive small-bowel obstruction: a propensity score-matched analysis.  
Scand J Surg. 2017;106(1):28-33
- 93. Chopra R , McVay C , Nemir A , Philips E,et al**  
laparoscopic lysis of Adhesions.  
Am Surg2003;69:966-968.
- 94. Wullstein C, Gross E.**  
Laparoscopic compared with conventional treatment of acute adhesive small bowel obstruction.  
Br J Surg. 2003;90(9):1147-51.
- 95. Li MZ, Lian L, Xiao LB, Wu WH, He YL, Song XM.**  
Laparoscopic versus open adhesiolysis in patients with adhesive small bowel obstruction: a systematic review and meta-analysis.  
Am J Surg 2012; 204: 779-786 [PMID: 22794708 DOI: 10.1016/ j.amjsurg.2012.03.005
- 96. Farinella E, Cirocchi R, La MF, Morelli U, Cattorini L, Delmonaco P, et al.**  
Feasibility of laparoscopy for small bowel obstruction.  
World J Emerg Surg. 2009;4:3
- 97. Michel cotton**  
Bowel obstruction  
Chapter 12 ,Primary surgery Volume one :non-trauma , page 210-240
- 98. DeWilde RL et Trew G.**  
Postoperative abdominal adhesions and their prevention in gynaecological surgery. Expert consensus position. Part 2 – steps to reduce adhesions Gynecol Surg. 2007; doi: 10.1007/s10397-007-0338.
- 99. Trésallet C, Royer B, Menegaux F.**  
Occlusion aigues du grêle de l'adulte.

EMC Techniques chirurgicales–appareil digestif 2010 ; 53:430–40.

**100. Barkan, H., S. Webster, and S. Ozeran.**

Factors predicting the recurrence of adhesive small-bowel obstruction.

Am J Surg, 1995. 170(4): p. 361-5.

**101. Landercasper, J., et al.**

Long-term outcome after hospitalization for small-bowel obstruction.

Arch Surg, 1993. 128(7): p. 765-70; discussion 770-1.

**102. Duron JJ, Silva NJ, du Montcel ST, Berger A, Muscari F, Hennet H, Veyrieres M, Hay JM.**

Adhesive postoperative small bowel obstruction: incidence and risk factors of recurrence after surgical treatment: a multicenter prospective study.

Ann Surg 2006; 244: 750–757

**103. Case courtesy of Dr Maryam Saif AlAli**

Radiopaedia.org, rID: 43438 <https://radiopaedia.org/cases/small-bowel-obstruction-19?lang=us>

**104. Drake, Vogl & Mitchell**

Gray's Anatomy for Students, 2nd Edition 2010. Chapter 4 : abdomen ,regional anatomy  
268–382p

**105. Anne M. Gilroy and Brian R. MacPherson**

Atlas of Anatomy – Third Edition. , Abdomen , 159– 244p

**106. Sherwood, Lauralee.**

Human Physiology : from Cells to Systems 7<sup>th</sup> edition. , chapter 16 : digestive system , small intestine 621–633 p .

**107. Kenneth Saladin**

Anatomy & Physiology: The Unity of Form and Function 8th Edition , Chapter 25 : digestive system , 944 – 985 p

**108. Allison H.MaciverMichaelMcCallA.M.James Shapiro .**

Intra–abdominal adhesions: Cellular mechanisms and strategies for prevention

International Journal of SurgeryVolume 9, Issue 8, 2011, Pages 589–594

**109. Taufiek Konrad Rajab et al**

Practical implications of postoperative adhesions for preoperative consent and operative technique

International Journal of Surgery ,Volume 11, Issue 9, November 2013, Pages 753–756

**110. Becker JM, Dayton MT, Fazio VW, et al.**

Prevention of postoperative abdominal adhesions by a sodium hyaluronate-based bioresorbable membrane: a prospective, randomized, double-blind multicenter study. *J Am Coll Surg.* 1996;183:297–306.

**111. Catena F, Ansaloni L, Di Saverio S, Pinna AD; World Society of Emergency Surgery et al.**

P.O.P.A. study: prevention of postoperative abdominal adhesions by icodextrin 4% solution after laparotomy for adhesive small bowel obstruction. A prospective randomized controlled trial. *J Gastrointest Surg.* 2012;16:382–388.

## قسم الطبيب

اقسمُ باللهِ العَظِيمِ

أن أراقبَ اللهَ في مِهْنَتِي.

وأن أصونَ حياةَ الإنسانِ في كافَّةِ أطوارها في كلِّ الظروفِ والأحوالِ

بإذلا وسعي في استنقاذها من الهلاكِ والمرَضِ والألمِ والقلقِ.

وأن أحفظَ للنَّاسِ كرامَتَهُم، وأستُرَّ عَوْرَتَهُم، وأكتُمَ سِرَّهُم.

وأن أكونَ على الدوامِ من وسائلِ رحمةِ اللهِ،

بإذلا رِعايتي للطبِّيةِ للقريبِ والبعيدِ، للصالحِ والطالحِ، والصديقِ والعدوِّ.

وأن أثابرَ على طلبِ العلمِ، أُسخره لنفعِ الإنسانِ .. لا لأذاهِ.

وأن أوقِّرَ مَنْ عَلَّمَنِي، وأُعَلِّمَ مَنْ يَصغرنِي، وأكونَ أبا لِكُلِّ زَميلٍ في المِهْنَةِ الطَّبِّيةِ

مُتعاونينَ على البرِّ والتقوى.

وأن تكونَ حياتي مُصدِّقَ إيمانِي في سِرِّي وَعَلائِيتِي ،

نَقِيَّةً مِمَّا يَشِينُهَا تَجَاهَ اللهِ وَرَسولِهِ وَالْمُؤمِنِينَ.

واللهِ على ما أقولُ شهيدٌ



كلية الطب  
والصيدلة - مراكش  
FACULTÉ DE MÉDECINE  
ET DE PHARMACIE - MARRAKECH

سنة 2019 أطروحة رقم 234

## الانسداد المعوي بالالتصاقات عند الاطفال

### الأطروحة

قدمت ونوقشت علانية يوم 2019/10/01

من طرف

السيد يونس الخدير

المزداد في 1993/09/27 بقصبة تادلة

لنيل شهادة الدكتوراه في الطب

الكلمات الأساسية :

الالتصاقات - الانسداد المعوي بالالتصاقات - الاطفال - المعالجة التحفظية  
- العامل التبايني الذواب بالماء

### اللجنة

الرئيس	السيد س. يونس
المشرف	السيد م. اولاد الصياد
الحكام	السيد ا.ا. كاملي
	السيد ه. جلال
	السيد م. بو الروس
	السيد أستاذ في طب الاطفال